

UNIFORMLY DISTRIBUTED SEQUENCES OF p -ADIC INTEGERS, II

VLADIMIR ANASHIN

ABSTRACT. The paper describes ergodic (with respect to the Haar measure) functions in the class of all functions, which are defined on (and take values in) the ring $\mathbb{Z}_p$ of p -adic integers, and which satisfy (at least, locally) Lipschitz condition with coefficient 1. Equiprobable (in particular, measure-preserving) functions of this class are described also. In some cases (and especially for $p = 2$) the descriptions are given by explicit formulae. Some of the results may be viewed as descriptions of ergodic isometric dynamical systems on the p -adic unit disk. The study is motivated by the problem of pseudorandom number generation for computer simulation and cryptography. From this view the paper describes nonlinear congruential pseudorandom generators modulo m which produce strictly periodic uniformly distributed sequences modulo m with maximal possible period length (i.e., exactly m). Both the state change function and the output function of these generators could be, e.g., meromorphic on $\mathbb{Z}_p$ functions (in particular, polynomials with rational, but not necessarily integer coefficients), or compositions of arithmetical operations (like addition, multiplication, exponentiation, raising to integer powers, including negative ones) with standard computer operations, such as bitwise logical operations (e.g., XOR, OR, AND, NEG, etc.). The linear complexity of the produced sequences is also studied.

1. INTRODUCTION.

A number of applications in computer simulation, numerical analysis (especially Quasi Monte Carlo) and cryptography demand regular methods to generate successively a uniformly distributed sequence. The corresponding literature is so vast that we could not even mention here the most important monographs in the area. We refer only [2], where a reader could find a rather substantial survey of relevant methods as well as a comprehensive bibliography. The major part of these methods are certain recursive procedures, which may be viewed also as automata. The latter are commonly referred as pseudorandom (or quasirandom) generators.

The typical one is the so-called linear congruential generator, which has been developed more than half a century ago. It produces a sequence $\{x_n : n = 0, 1, 2, \dots\}$ over a set $\{0, 1, \dots, m-1\}$ (the latter is commonly treated as the residue class ring $\mathbb{Z}/m$ of the ring $\mathbb{Z}$ of rational integers modulo natural $m > 1$), which is a first order recurrence sequence, defined by $x_{n+1} \equiv a + bx_n \pmod{m}$ with integer rationals a, b . The sequence is uniformly distributed iff it is purely periodic with period length m . The latter condition implies that *each* element of $\mathbb{Z}/m$ occurs at

Key words and phrases. Uniformly distributed sequence; p -adic integer; non-Archimedean dynamical system; ergodic function; equiprobable function; measure-preserving function; transitive polynomial; bijective polynomial; permutation polynomial; pseudorandom number generator; non-linear congruential generator; linear complexity.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

the period *exactly once*; and vice versa. The necessary and sufficient conditions a and b must (for a given m) satisfy to provide the maximal period length (i.e., m) of the produced sequence, are well known –see [2, section 3.2.1.2, theorem A].

The undoubtful advantage of linear congruential generators is the simplicity (especially for $m = 2^k$) of their program implementations. One of the key reasons of their disadvantages (e.g., lack of statistical quality of the produced sequences, for certain applications) is their linearity. For instance, as the state change function $f(x) = a + bx$ of the generator is a polynomial of degree 1, the produced sequence has linear complexity 2 over the ring $\mathbb{Z}/m$, i.e., it is a linear recurrence sequence of order 2 over $\mathbb{Z}/m$ (defined by $x_{n+2} \equiv (1+b)x_{n+1} - bx_n \pmod{m}$). Hence, *for each* m the points $(\frac{x_{n+2}}{m}, \frac{x_{n+1}}{m}, \frac{x_n}{m})$ fall into the parallel planes $c + X - (1+b)Y + bZ$ ($c \in \mathbb{Z}$), which intersect the unit cube of Euclidean space. The well known result due to George Marsaglia [7] states that similar effect also holds in higher dimensions > 3 : all the points fall into the relatively small number of parallel hyperplanes (rather than fill this cube more or less uniformly), and the reason is again that $\deg f = 1$.

During the past decades these considerations stimulated the developement of various alternatives to linear congruential generators. The significant part of these are *nonlinear* congruential generators with state change function f being either a polynomial over $\mathbb{Z}$ of degree > 1 , (in particular, quadratic [2], or of higher degree [15]), or some non-polynomial transformations, which gave rise to exponential generators (with $f(x) = a^{g(x)}$), or to so-called inversive generators, involving raising to negative powers (for the survey of different generators we again refer to [2]). Very often some authors seem to be more concerned with the linear complexity of the produced sequence, then with its uniform distribution, admitting non-maximal period length, i.e., they admit state change functions f , for which the sequence never reaches the period length m , and hence, in a strict sence, is not uniformly distributed in $\mathbb{Z}/m$. In such cases the authors have not only to estimate possible period lengths, but also to choose the initial state (the *seed*) x_0 of the generator according to certain (sometimes, sophisticated) procedures, which are to assure logging on the sufficiently long period, rather then to choose the seed at random.

Increasing the degree of a polynomial (as well as the use in the composition other arithmetical operations like exponentiation or taking an inverse) is also has to be paid for by certain rise of complexity of program implementation. The most disappointing here is that the statistical quality and complexity of the program implementation of the generators often occur to be in inverse dependence: the better the quality the slower the performance; fast generators sometimes demonstrate lack of quality.

So it is still important to find new classes of functions $f: \mathbb{Z}/m \rightarrow \mathbb{Z}/m$, for which the corresponding generators,

- (1) firstly, achieve maximal possible (i.e., exactly m) period length of the recurrence sequence defined by the relation $x_{n+1} \equiv f(x_n) \pmod{m}$, hence producing uniformly distributed sequence in $\mathbb{Z}/m$ (we refer such transformations f as *transitive modulo m*);
- (2) secondly, guarantee the sufficiently large linear complexity of the produced sequence over $\mathbb{Z}/m$, i.e., absence of ‘short’ (in some definite sence) linear dependencies of the form $\sum_{i=0}^{r-1} c_i x_{n+i} \equiv 0 \pmod{m}$ ($n = 0, 1, 2, \dots$) among the members of the sequence;

- (3) and, thirdly, basically are ‘easy-to-implement’, namely, are ‘flexible’, i.e., have certain (critical to the performance) parameters, varying which it is possible to gain speed without losing quality.

The paper presents wide classes of transformations f which to some extent satisfy these conditions.

At the first turn we obtain transitivity modulo m conditions for functions, which could be implemented as compositions of arithmetical operations (addition and multiplication of integers), as well as of standard computer ones, like bitwise logical operations, shifts, masking, etc. These compositions might involve as well exponentiation and taking a multiplicative inverse, hence, raising to negative powers (see 4.9, 4.11, 2.5) and/or OR, XOR, AND, etc., see 2.5, 2.8.

In particular, we describe wide classes of transitive modulo m functions which could be expressed as integer-valued polynomials with rational (and not necessarily integer) coefficients (see 4.7), as well as by analytic functions (4.11, 4.9, 2.5) or meromorphic (in particular, rational) functions (4.9, 4.11, 4.12). These conditions are easy-to-verify, and with the use of them the various explicit formulae for transitive modulo m transformations could be (and are) obtained – see e.g. 2.3, 2.4, also 2.5–2.8 (as well 4.11, 4.12) together with 2.1, and other examples here and there in the paper.

To illustrate, we start with some of these examples: theorem 2.7 together with lemma 4.11 imply that each transformation f of the form

$$f(x) = 1 + x + 2(g(x+1) - g(x))$$

is transitive modulo $m = 2^k$ for all $k = 1, 2, \dots$ and for arbitrary composition g of

- (1) arithmetical operations — an addition $(y, z) \mapsto y + z$, a multiplication $(y, z) \mapsto yz$, an exponentiation $(y, z) \mapsto (1 + 2y)^z$ (in particular, taking an inverse $y \mapsto (1 + 2y)^{-1}$),
- (2) bitwise logical operations — such as conjunction $(y, z) \mapsto y \text{ AND } z$, disjunction $(y, z) \mapsto y \text{ OR } z$, exclusive ‘or’ $(y, z) \mapsto y \text{ XOR } z$, negation $z \mapsto \text{NEG } z$, etc.,
- (3) machine operations (which could be derived from the bitwise logical ones) — an s -step shift towards most significant bits $z \mapsto 2^s z$, masking $z \mapsto z \text{ AND } M$, M being a mask, ‘reduction modulo 2^s ’, i.e., a truncation of the most significant bits $z \mapsto z \bmod 2^s = z \text{ AND } (2^s - 1)$, and some others.

We assume here that all the operands are non-negative integer rationals which are represented as base 2 expansions; so, for instance, $2 = 1 \text{ XOR } 3 = 2 \text{ AND } 7 \equiv \text{NEG } 13 \pmod{8}$, $3^{-1} \equiv 11 \equiv -5 \pmod{16}$, $3^{-\frac{1}{3}} \equiv 3^{11} \equiv 3^{-5} \equiv 11 \pmod{16}$, etc. Up to this agreement the functions g and f are correctly defined on $\mathbb{Z}/m$, the efficiency of their program implementation depends only on the number of ‘fast’ and ‘slow’ operations in the composition g and hence one may vary it in wide range to achieve the desired performance.

We emphasize, in the example just mentioned the transitivity modulo $m = 2^k$ does not depend neither on k nor on actual form of the composition g — both for $g(x) = x \text{ XOR } (2x + 1)$ and for

$$g(x) = \left(1 + 2 \frac{x \text{ AND } x^2 + x^3 \text{ OR } x^4}{3 + 4(5 + 6x^5)^{x^6} \text{ XOR } x^7} \right)^{7 + \frac{8x^8}{9 + 10x^9}}$$

3

the sequence $\{x_n\}$ defined by the recurrence relation $x_{n+1} \equiv 1 + x_n + 2(g(x_n + 1) - g(x_n)) \pmod{2^k}$ is uniformly distributed in $\mathbb{Z}/2^k$ for each $k = 1, 2, 3, \dots$. Actually, this sequence is strictly periodic with period length 2^k , and *each* element of $\{0, 1, \dots, 2^k - 1\}$ occurs at the period *exactly once*.

Similar assertions also hold for arbitrary composite m : e.g., 4.11 and 4.12 imply that the transformation

$$f(x) = 1 + x + \pi(m)^2 u(x)(1 + \pi(m)v(x))^{w(x)}$$

with $\pi(m)$ being a product of all prime factors of m , is transitive modulo m for arbitrary polynomials $u(x), v(x), w(x) \in \mathbb{Z}[x]$ over $\mathbb{Z}$. A variety of results of such kind may be obtained in much more general situation for integer-valued polynomials with rational (not necessarily integer) coefficients by applying the techniques of section 4.

Note that this example also demonstrates how by minor changes of the recurrence relation one may achieve the transitivity of both inversive generator (for which $f(x) = a + bx^{-1}$ or $f(x) = (a + bx)^{-1}$) and exponential generator (with $f(x) = a^x$): for $w(x) = \text{const} = -1$ the introduced generator is of inversive type, for $v(x) = \text{const} \neq 0$ it is of exponential type.

As for linear dependencies $\sum_{i=0}^{r-1} c_i x_{n+i} \equiv 0 \pmod{m}$ ($n = 0, 1, 2, \dots$) of fixed length r in produced sequences $\{x_n \equiv f(x_{n-1}) \pmod{m} : n = 1, 2, \dots\}$, one may say that from this view among all congruential generators linear ones are rather exceptions than the law. For instance, if $f: \mathbb{Z} \rightarrow \mathbb{Z}$ is represented by a transitive modulo some prime power $m = p^k$ ($k \geq 3$) polynomial of degree ≥ 2 with integer rational coefficients, no such dependencies with r and c_i not depending on k do exist. Moreover, in this case the minimal order of linear recurrence sequence over $\mathbb{Z}/p^k$, which represents the produced sequence, tends to infinity together with k (in fact, much more general result holds — see 5.1–5.4 for exact statements).

The paper also studies equiprobable modulo m functions, i.e., mappings F of the s th Cartesian power $(\mathbb{Z}/m)^{(s)}$ onto the t th Cartesian power $(\mathbb{Z}/m)^{(t)}$ of the ring $\mathbb{Z}/m$, ($s \geq t$), such that all preimages of all elements are of the same cardinality. In particular, for $s = t$ equiprobable modulo m functions are bijections of the corresponding rings and throughout the paper are referred as bijective modulo m functions. A very particular case of the equiprobable modulo m functions studied here are so-called permutation polynomials modulo m , the latter being polynomials over $\mathbb{Z}$ which induce bijections of the ring $\mathbb{Z}/m$ onto itself. The results of the paper concerning equiprobability modulo m generalize known [8] results on permutation polynomials to much wider classes of functions. The study was motivated by the observation that application of equiprobable modulo m functions as output functions to uniformly distributed in $\mathbb{Z}/M$ periodic sequences with period length M leads to new uniformly distributed in $\mathbb{Z}/N$ (with $N|M$) sequences of the same period length M . In other words, each element of $\mathbb{Z}/N$ occurs at the period of such sequence the same number of times (but not necessarily once). Hashing with equiprobable modulo m functions the sequences, generated by already introduced methods, seems to be useful to design secure stream ciphers. Yet this will be an issue of the forthcoming paper and is out of the scope of the present one.

Note that proofs of our basic assertions imply p -adic techniques. The problems stated above are firstly restated in these terms. Actually the paper studies ergodic with respect to the Haar measure (as well as preserving this measure or equiprobable

with respect to it) functions, which are defined on (and which take values in) the space $\mathbb{Z}_p$ of all p -adic integers, and which are non-expanding functions, i.e., satisfy Lipschitz condition with coefficient 1. From this view the results of the paper could be of interest for non-Archimedean dynamical systems theory: a number of statements could be easily interpreted as descriptions of ergodic dynamical systems with discrete time and with $\mathbb{Z}_p$ as a phase space.

The paper continues the study started in [11]: here we prove some results announced in [11, 12, 14, 17] and establish new ones. Moving towards exact statements, for reader's convenience we recall some facts from the p -adic analysis and the theory of uniform distribution of sequences, following [6], [3] and [2]; we recall some necessary results, definitions and notations from [11] as well.

Here and after let p be a prime number. Consider a canonic representation $z = z_0 + z_1p + z_2p^2 + \dots$ of p -adic integer $z \neq 0$, where $z_j \in \{0, 1, \dots, p-1\}$ ($j = 0, 1, 2, \dots$); we denote $\text{ord}_p z = \min\{j : z_j \neq 0\}$ the exponent of a maximal power of p which is a factor of z . By definition, $\|z\|_p = p^{-\text{ord}_p z}$ is p -adic norm of z , $\|0\|_p = 0$. The valuation $\|\cdot\|_p$ could be expanded to the whole field $\mathbb{Q}_p$ of p -adic numbers (which is a quotient field of the ring $\mathbb{Z}_p$ of p -adic integers) in a standard way; so this valuation induces on $\mathbb{Q}_p$ a distance $d_p(u, v) = \|u - v\|_p$, with $\mathbb{Q}_p$ being a completion of the space $\mathbb{Q}$ of all rationals with respect to this distance. Note that often they use another terminology, where a distance is called a *metric*, a p -adic norm is called a *p -adic value*, whereas the term ' p -adic valuation' is reserved for ord_p . However, throughout the paper we mainly use the terminology of [3], with the only exception, speaking of ' p -adic norms' instead of ' p -adic values'.

The ring $\mathbb{Z}_p = \{u \in \mathbb{Q}_p : \|u\|_p \leq 1\}$ is compact in the space $\mathbb{Q}_p$, being a closure of the set $\mathbb{N}_0 = \{0, 1, 2, \dots\}$. Hence, $\mathbb{Z}_p$ is a separable compact metric space. The set of all cosets with respect to all ideals of the ring $\mathbb{Z}_p$ forms a base of the corresponding topology. Each coset $a + p^k\mathbb{Z}_p$ ($a \in \mathbb{Z}_p$, $k = 0, 1, 2, \dots$) is an open (and simultaneously closed) ball of radius p^{-k} .

There exists a natural measure μ on $\mathbb{Z}_p$: putting $\mu(a + p^k\mathbb{Z}_p) = p^{-k}$, we then expand μ to the corresponding σ -ring generated by all compact subsets of $\mathbb{Z}_p$ (these compact subsets are exactly all closed subsets of $\mathbb{Z}_p$). So we define uniquely a measure on $\mathbb{Z}_p$, which is non-negative σ -additive regular normalized Borel and Haar measure in this case. Thus, μ is a natural probability measure on $\mathbb{Z}_p$. The probability measure on n -dimensional space $\mathbb{Z}_p^{(n)}$ could be defined in a similar way as a corresponding normalized Haar measure.

Now let $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be a function, which preserves all congruences of the ring $\mathbb{Z}_p$, i.e., $a\theta b$ implies $f(a)\theta f(b)$ for each congruence θ and all $a, b \in \mathbb{Z}_p$. As each nontrivial congruence of the ring $\mathbb{Z}_p$ is an equivalence modulo certain ideal $p^k\mathbb{Z}_p$ (we denote this congruence via $\cdot \equiv \cdot \pmod{p^k}$), it can be easily shown that the function f preserves all the congruences of the ring $\mathbb{Z}_p$ iff it satisfies Lipschitz condition with coefficient 1: $\|f(x) - f(y)\|_p \leq \|x - y\|_p$. Function preserving all congruences of a universal algebra is called *compatible*; we will use this term instead of the term '*conservative*' of [11], since the latter in numerous papers on algebraic systems has another meaning, see [8, p. 45].

The class of all compatible functions on $\mathbb{Z}_p$ is rather wide: it contains all functions represented by polynomials with rational integer or p -adic integer coefficients, integer-valued analytic on $\mathbb{Z}_p$ functions, as well as integer-valued and meromorphic (in particular, rational) on $\mathbb{Z}_p$ functions with denominators equivalent to 0 modulo

p at no point of $\mathbb{Z}_p$. Some other examples will be introduced further in the paper.

Recall that a function, which is defined on some field F , and which takes values there, is called *integer-valued* iff all its values are integers of F providing arguments take integer values in F . Further we study integer-valued functions on the field $\mathbb{Q}_p$; hence they map $\mathbb{Z}_p$ into $\mathbb{Z}_p$. In particular, we study integer-valued functions on $\mathbb{Q}$. A polynomial over a field F is called integer-valued iff it induces an integer-valued function on F . Note that integer-valued function f on F defines on the ring Z of all integers of F a function $f|_Z : Z \rightarrow Z$, which is not necessarily compatible on Z , i.e., does not necessarily preserve all congruences of Z ; yet, if $f|_Z$ is compatible as a function on Z , then (in cases which do not lead to misunderstanding) we also call f compatible. Moreover, if a compatible integer-valued function f could be defined by a polynomial over F , we call compatible the corresponding polynomial too.

Note that the notion of compatible integer-valued function could be naturally expanded to the multivariate case — a valuation (and hence, a distance) on the space $\mathbb{Z}_p$ induces a (pseudo)-valuation (hence, a distance) on the n -dimensional space $\mathbb{Z}_p^{(n)}$ in a standard manner: for $\mathbf{u} = (u_1, \dots, u_n) \in \mathbb{Z}_p^{(n)}$ we assume $\|\mathbf{u}\|_p = \max\{\|u_i\|_p : i = 1, 2, \dots, n\}$. So, the function

$$F = (f_1, \dots, f_m) : \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$$

is compatible iff it satisfies Lipschitz condition with coefficient 1. In particular, all compatible on $\mathbb{Z}_p$ functions are continuous as functions of p -adic variables.

This obvious conclusion is important for applications. Each machine word, i.e., a word of some finite length in the alphabet $\{0, 1\}$, could be treated as a 2-base expansion of a non-negative integer rational. Then all the above mentioned bitwise logical operations and machine operations could be naturally continued to the set $\mathbb{Z}_2$ of all 2-adic integers in their canonic representations. Moreover, the above mentioned arithmetical operations could be continued to $\mathbb{Z}_2$ either. It could be easily demonstrated that all these operations (to be more precise, their uniquely defined continuations to $\mathbb{Z}_2$) and all their compositions are compatible (hence, continuous) integer-valued functions on $\mathbb{Z}_2$: for exponentiation $(y, z) \mapsto (1 + 2y)^z$, and, in particular, for the inversion $y \mapsto (1 + 2y)^{-1}$ see 4.11, for the rest the assertion follows immediately from the corresponding definitions. We note here that an m -step shift towards less significant bits (i.e., the operation $\lfloor \frac{\cdot}{2^m} \rfloor$ of ‘integer division’, a division succeeded by a truncation of the fractional part of the quotient) is *not* compatible, yet continuous, integer-valued function on $\mathbb{Z}_2$ (hence the results of the paper remain valid for compositions including the latter operation either, providing the whole composition is compatible).

These considerations give an opportunity to apply, while studying compositions of the above mentioned operations, certain methods of non-Archimedean (p -adic) analysis. Certainly, these techniques could be applied only to problems which are stated in appropriate terms (measures, distances, limits, derivatives, etc.).

It turns out that some properties of functions, which traditionally have been treated as discrete mathematics issues, could be restated in these terms. We have already introduced one of such properties, namely, compatibility. It worth a brief notice in this connection that so-called ‘determinate functions on superwords’ of automata theory (which are functions defined on infinite sequences of $\{0, 1\}$) after natural identification of superwords with elements of $\mathbb{Z}_2$ could be considered as compatible functions on $\mathbb{Z}_2$.

There exist other properties which could be restated in such manner. We consider a property of a compatible function $F = (f_1, \dots, f_m): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$ to be *equiprobable modulo p^k* . The latter by definition means that the function F induces on the n th Cartesian power $(\mathbb{Z}/p^k)^{(n)}$ of $\mathbb{Z}/p^k$ an equiprobable function $\bar{F} = (\bar{f}_1, \dots, \bar{f}_m): (\mathbb{Z}/p^k)^{(n)} \rightarrow (\mathbb{Z}/p^k)^{(m)}$, i.e., each point of $(\mathbb{Z}/p^k)^{(m)}$ has the same number of F -preimages in $(\mathbb{Z}/p^k)^{(n)}$. In particular, for $m = n$ equiprobable modulo p^k functions are exactly *bijective modulo p^k* functions. We consider also an important (especially for pseudorandom generation) property of a bijective modulo p^k function F to be *transitive modulo p^k* , which means that F induces on $(\mathbb{Z}/p^k)^{(n)}$ a single cycle permutation. Note that while defining notions of equiprobability, bijectivity or transitivity of a function F modulo p^k , we have assumed the compatibility of F .

A value of induced function $\bar{f}_i(x)$ in the ring $\mathbb{Z}/p^k$ is, by definition, $f_i(x) \bmod p^k$, the least non-negative residue modulo p^k of $f_i(x)$, i.e., $f_i(x) \bmod p^k = \alpha \in \{0, 1, \dots, p^k - 1\}$, with $\|f_i(x) - \alpha\|_p \leq p^{-k}$. In view of compatibility of the function f_i , the value of the function $\bar{f}_i(x)$ does not depend on choice of the representative x in a coset of the ring $\mathbb{Z}_p^{(n)}$ with respect to the ideal $(p^k \mathbb{Z}_p)^{(n)}$; hence, the function F correctly defines on $(\mathbb{Z}/p^k)^{(n)}$ a function $F \bmod p^k = (f_1(x) \bmod p^k, \dots, f_m(x) \bmod p^k)$, which takes values in $(\mathbb{Z}/p^k)^{(m)}$. Throughout the paper the latter function is denoted via $F \bmod p^k$, or via $\bar{F}$, when it does not lead to misunderstanding.

Now recall some definitions of the theory of measurable functions (cf. [1]). Let S and T be spaces with nonnegative normalized measures μ and τ , respectively, and let $f: S \rightarrow T$ be a measurable function, i.e., each full f -preimage $f^{-1}(U)$ of $U \subseteq T$ is μ -measurable for each τ -measurable U .

We say that the function f is (μ, τ) -*proportional*, iff for each pair of τ -measurable subsets $U, V \subseteq T$ the equality $\tau(U) = \tau(V)$ implies the equality $\mu(f^{-1}(U)) = \mu(f^{-1}(V))$. In case both μ, τ are probability measures (e.g., are properly normalized Haar measures), then f is called (μ, τ) -*equiprobable* (or *equiprobable with respect to μ and τ*) iff $\mu(f^{-1}(U)) = \tau(U)$ for each τ -measurable $U \subseteq T$. For $S = T$ and $\mu = \tau$ we say that f *preserves measure μ* , iff $\mu(f^{-1}(U)) = \mu(U)$ holds for each μ -measurable U . Finally, if f preserves measure μ , and for μ -measurable subset U the equality $f^{-1}(U) = U$ implies that either $\mu(U) = 0$, or $\mu(U) = 1$, we say that f is μ -*ergodic* (or *ergodic with respect to μ*).

Note that in metric theory instead of terms ‘measure-preserving function’ or ‘equiprobable function’ they often use terms ‘metric endomorphism’ and ‘metric homomorphism’, and in dynamical systems theory they sometimes speak about ‘metric transitivity’ instead of ergodicity. Since throughout the paper we deal with the only measure, the properly normalized Haar measure, we omit mentioning this measure, so preserving the Haar measure, equiprobable (accordingly, ergodic) with respect to the Haar measure functions are referred as *measure-preserving, equiprobable* (or, accordingly, *ergodic*).

The following theorem holds:

1.1 Theorem. *A compatible function $F: \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$ is equiprobable (respectively, measure-preserving or ergodic) iff it is equiprobable (respectively, is bijective or transitive) modulo p^k for all $k = 1, 2, \dots$. A compatible and measure-preserving function F is bijective (consequently, is a metric automorphism); moreover, it is*

an isometry of the space $\mathbb{Z}_p^{(n)}$.

Note that further throughout the paper while proving ergodicity (equiprobability) of a compatible function with respect to the Haar measure we actually prove its transitivity (equiprobability) each modulo p^k , $k = 1, 2, \dots$, i.e., directly establish the properties we are interested in view of the problems mentioned above. That is why we omit the proof of this theorem 1.1: it is not related directly to the aims of this paper. Nevertheless throughout the paper we use the relevant terminology (e.g., we commonly speak of ‘ergodicity’ instead of ‘transitivity modulo p^k for all $k = 1, 2, \dots$ ’, etc.)

In connection with theorem 1.1 it is worth noticing, however, that the results of the paper related to description of measure-preserving or ergodic functions may be treated as description of non-Archimedean (i.e., ultrametric) dynamical systems $(\mathbb{Z}_p^{(n)}, F)$ with phase space $\mathbb{Z}_p^{(n)}$, discrete time, and with nonexpanding F (i.e. for each pair of points $\mathbf{a}, \mathbf{b}$ a distance between their F -images $F(\mathbf{a})$ and $F(\mathbf{b})$ does not exceed a distance between these points). In this sense theorem 2.2, for instance, might be considered as a complete description (in terms of explicit formulae) of ergodic dynamical systems of the above mentioned kind when $p = 2$ and $n = 1$; together with theorem 3.11 it gives full description of twice integer-valued (i.e., having everywhere integer-valued derivative) ergodic dynamical systems. These themes, however, are not covered by this paper and will be considered in forthcoming one.

Returning to the leading theme of the paper we note that for a wide class of compatible functions, which are in some (properly defined in section 3) sense generalizations of uniformly differentiable on $\mathbb{Z}_p$ functions, the bijectivity modulo p^k of a function for a *certain* k is equivalent to the property of being measure-preserving; the latter is equivalent to its bijectivity modulo p^k for *all* $k = 1, 2, 3, \dots$. The property of being transitive modulo p^k for a *certain* k turned out to be equivalent to the ergodicity of a function; the latter implies that the function is transitive modulo p^k for *all* $k = 1, 2, 3, \dots$. Finally, the equiprobability of a function modulo p^k for a *certain* k implies its equiprobability with respect to the Haar measure; the latter property is equivalent to equiprobability modulo p^k for *all* $k = 1, 2, 3, \dots$. The results of this kind are proved in section 3.

These results demonstrate the same remarkable effect originally enlightened by Hensel lemma: the Hensel lift, that is, a situation when a behavior of a function modulo p^{k_0} for a certain k_0 controls its behavior modulo p^k for all $k = k_0 + 1, k_0 + 2, \dots$ and on the whole space $\mathbb{Z}_p$. This effect have been already observed while studying transitivity of some transformations. For instance, the necessary and sufficient conditions for the polynomial $f(x) = a + bx$ with integer rational a, b (see e.g., [2; 3.2.1.2, theorem A]) could be restated as follows: a polynomial $a + bx$ is transitive modulo p^k for some (that is, for all) $k \geq 2$ iff it is transitive modulo p for odd p or, respectively, modulo p^2 for $p = 2$. The general criterion for the transitivity modulo p^k of the polynomial f of arbitrary degree over integer rationals [15] demonstrates this effect either: for $p \neq 2, 3$ a polynomial f is transitive modulo p^k , $k \geq 3$, iff it is transitive modulo p^2 ; respectively, for $p = 2$ or $p = 3$ — iff it is transitive modulo p^3 . Note by the way that the latter result holds for a much wider class of functions, even not necessarily analytic (see 4.9–4.10).

The results of section 3 show that Hensel lift of such properties as bijectivity or transitivity modulo p^k is basically due to the specific character of p -adic distance

and holds for various rather wide classes of functions. The values of k_0 from which the lift starts are estimated in section 4.

The results of this kind are useful if for a given f one has to establish whether it shares some property (e.g., transitivity or bijectivity) modulo p^k for a definite rather large k , for which direct verification is not accessible. However, if one needs to construct out of prescribed operations a certain function, which is to be transitive or bijective modulo p^k , then explicit formulae are more convenient. Such formulae for bijective modulo 2^k polynomials over $\mathbb{Z}$ were obtained in [13], for transitive modulo 2^k polynomials over $\mathbb{Z}$ — in [15]. Explicit formulae for ergodic or measure-preserving compatible functions (in particular, for compatible integer-valued polynomials over $\mathbb{Q}$), which are defined on (and take values in) $\mathbb{Z}_2$ were obtained in [11]. The current paper presents explicit formulae for compatible ergodic (or measure-preserving) functions on $\mathbb{Z}_p$ for odd p — see the next section.

2. EXPLICIT FORMULAE

Recall (see [3]) that each function $f: \mathbb{N}_0 \rightarrow \mathbb{Z}_p$ (or, respectively, $f: \mathbb{N}_0 \rightarrow \mathbb{Z}$) admits one and only one representation in the form of so-called *interpolation series*

$$f(x) = \sum_{i=0}^{\infty} a_i \binom{x}{i}, \quad (\diamond)$$

where $\binom{x}{i} = \frac{x(x-1)\cdots(x-i+1)}{i!}$ for $i = 1, 2, \dots$, and $\binom{x}{0} = 1$; $a_i \in \mathbb{Z}_p$ (respectively, $a_i \in \mathbb{Z}$), $i = 0, 1, 2, \dots$.

If f is uniformly continuous on $\mathbb{N}_0$ with respect to p -adic distance, it can be uniquely continued to the uniformly continuous function on $\mathbb{Z}_p$. Hence the interpolation series for f converges uniformly on $\mathbb{Z}_p$. The following is true: the series $f(x) = \sum_{i=0}^{\infty} a_i \binom{x}{i}$, ($a_i \in \mathbb{Q}_p$, $i = 0, 1, 2, \dots$) converges uniformly on $\mathbb{Z}_p$ iff $\lim_{i \rightarrow \infty}^p a_i = 0$, where $\lim$ is a limit with respect to p -adic distance; hence the uniformly convergent series defines a uniformly continuous function on $\mathbb{Z}_p$. The latter function is integer-valued iff $a_i \in \mathbb{Z}_p$ for all $i = 0, 1, 2, \dots$.

Further throughout this section we assume that the function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is uniformly continuous on $\mathbb{Z}_p$, and that it is represented by series $(\diamond)$. The following three criteria hold (see [11]):

2.1 Theorem. (See 4.3 of [11]; cf. [5]) *A function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is compatible iff*

$$a_i \equiv 0 \pmod{p^{\lfloor \log_p i \rfloor}}$$

for all $i = p, p+1, p+2, \dots$. (Here and after for a real α we denote $\lfloor \alpha \rfloor$ an integral part of α , i.e., the nearest to α integer rational not exceeding α .)

2.2 Theorem. (See 4.5 of [11]) *A function $f: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ is compatible and measure-preserving iff it could be represented as*

$$f(x) = c_0 + x + \sum_{i=1}^{\infty} c_i 2^{\lfloor \log_2 i \rfloor + 1} \binom{x}{i},$$

where $c_0, c_1, c_2, \dots \in \mathbb{Z}_2$.

2.3 Theorem. (See 4.7 of [11]) *A function $f: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ is compatible and ergodic iff it could be represented as*

$$f(x) = 1 + x + \sum_{i=0}^{\infty} c_i 2^{\lfloor \log_2(i+1) \rfloor + 1} \binom{x}{i},$$

where $c_0, c_1, c_2 \dots \in \mathbb{Z}_2$.

For an arbitrary prime p the necessity of conditions of theorems 2.2 and 2.3 does not hold, yet the sufficientness remains true. Namely, in this section we prove the following:

2.4 Theorem. *Let p be an odd prime. A function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$, which is represented in the form $\langle \diamond \rangle$, is compatible and measure-preserving if the following congruences hold simultaneously:*

$$\begin{aligned} a_1 &\not\equiv 0 \pmod{p}; \\ a_i &\equiv 0 \pmod{p^{\lfloor \log_p i \rfloor + 1}}, \quad (i = 2, 3, \dots). \end{aligned}$$

The function f is compatible and ergodic if the following congruences hold simultaneously:

$$\begin{aligned} a_0 &\not\equiv 0 \pmod{p}; \\ a_1 &\equiv 1 \pmod{p}; \\ a_i &\equiv 0 \pmod{p^{\lfloor \log_p(i+1) \rfloor + 1}}, \quad (i = 2, 3, \dots). \end{aligned}$$

For the proof of the theorem we will need two additional results which are of interest by their own.

2.5 Lemma. *Let p be an arbitrary prime, let $v: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be a compatible function, and let c, d be p -adic integers, with $c \not\equiv 0 \pmod{p}$. Then the function $g(x) = d + cx + pv(x)$ preserves measure, and the function $h(x) = c + x + p\Delta v(x)$ is ergodic. (Here and after Δ is a difference operator: $\Delta v(x) = v(x+1) - v(x)$. Note that both g and h are obviously compatible since they are compositions of compatible functions.)*

Proof of the lemma 2.5. Firstly by induction on l we show that g is bijective modulo p^l for all $l = 1, 2, 3, \dots$. The assumption is obviously true for $l = 1$.

Assume it is true for $l = 1, 2, \dots, k-1$. Prove that it holds for $l = k$ either. Let $g(a) \equiv g(b) \pmod{p^k}$ for some p -adic integers a, b . Then $a \equiv b \pmod{p^{k-1}}$ by the induction hypothesis. Hence $pv(a) \equiv pv(b) \pmod{p^k}$ since v is compatible. Further, the congruence $g(a) \equiv g(b) \pmod{p^k}$ implies that $ca + pv(a) \equiv cb + pv(b) \pmod{p^k}$, and consequently, $ca \equiv cb \pmod{p^k}$. Since $c \not\equiv 0 \pmod{p}$, the latter congruence implies that $a \equiv b \pmod{p^k}$, proving the first assertion of the lemma.

To prove the rest part of the statement we note firstly that the assertion just proven implies that h preserves measure. To prove the transitivity of h modulo p^k for all $k = 1, 2, 3, \dots$ we apply induction on k once again.

It is obvious that h is transitive modulo p . Assume that h is transitive modulo p^{k-1} . Then, since h induces a permutation on $\mathbb{Z}/p^k$ and since it is a compatible

function, we conclude that the length of each cycle of this permutation must be a multiple of p^{k-1} . So to prove this permutation is single cycle it is sufficient to prove that the function

$$h^{p^{k-1}}(x) = \underbrace{h(h \dots (h(x)) \dots)}_{p^{k-1}}$$

induces a single cycle permutation on the ideal (p^{k-1}) , generated by the element p^{k-1} of the ring $\mathbb{Z}/p^k$. In other words, it is sufficient to demonstrate that the function $\frac{1}{p^{k-1}}h^{p^{k-1}}(p^{k-1}x)$ is transitive modulo p .

Applying obvious direct calculations, we successively obtain that

$$h^1(x) = c + x + pv(x+1) - pv(x),$$

$$\dots \quad \dots \quad \dots$$

$$\begin{aligned} h^j(x) &= h(h^{j-1}(x)) = cj + h^{j-1}(x) + pv(h^{j-1}(x) + 1) - pv(h^{j-1}(x)) \\ &= cj + x + p \sum_{i=0}^{j-1} v(h^i(x) + 1) - p \sum_{i=0}^{j-1} v(h^i(x)), \end{aligned}$$

and henceforth. We recall that $h^0(x) = x$ by definition. So,

$$h^{p^{k-1}}(x) = cp^{k-1} + x + p \sum_{i=0}^{p^{k-1}-1} v(h^i(x) + 1) - p \sum_{i=0}^{p^{k-1}-1} v(h^i(x)). \quad (1)$$

Since h is transitive modulo p^{k-1} and compatible, we get now that

$$\sum_{i=0}^{p^{k-1}-1} v(h^i(x) + 1) \equiv \sum_{i=0}^{p^{k-1}-1} v(h^i(x)) \equiv \sum_{z=0}^{p^{k-1}-1} v(z) \pmod{p^{k-1}},$$

and (1) implies then $h^{p^{k-1}}(x) \equiv cp^{k-1} + x \pmod{p^k}$. But $c \not\equiv 0 \pmod{p}$, so we conclude that the function $cp^{k-1} + x$ induces on the ideal (p^{k-1}) a single cycle permutation, thus proving the lemma. $\square$

2.6 Corollary. *Under the assumptions of lemma 2.5, let p be an odd prime, and let $r \equiv 1 \pmod{p}$. Then the function $c + rx + p\Delta v(x)$ is compatible and ergodic.*

Proof of the corollary 2.6. We have that $r = 1 + ps$ for a suitable $s \in \mathbb{Z}_p$. Now, since p is odd, the function $s\binom{x}{2}$ is compatible; consequently, the function $v_1(x) = s\binom{x}{2} + v(x)$ is compatible either. Yet $\Delta v_1(x) = sx + \Delta v(x)$, and it is sufficient now to apply lemma 2.5 to finish the proof of the corollary. $\square$

Proof of the theorem 2.4. Note that according to 2.1 a compatible function $v(x)$ could be represented as

$$v(x) = a + \sum_{i=1}^{\infty} b_i p^{\lfloor \log_p i \rfloor} \binom{x}{i},$$

where $a, b_1, b_2, \dots \in \mathbb{Z}_p$. As $\lfloor \log_p i \rfloor = \lfloor \log_p(i+1) \rfloor$ for all $i = 1, 2, \dots$ with the exception of $i = p^t - 1$, ($t = 1, 2, 3, \dots$), and as

$$\Delta v(x) = \sum_{i=1}^{\infty} b_i p^{\lfloor \log_p i \rfloor} \binom{x}{i-1}, \quad (1)$$

we finish the proof of the theorem, applying 2.5 and 2.6. $\square$

For $p = 2$ the results just proven imply one more useful criterion of ergodicity of a function (or being measure-preserving).

2.7 Theorem. *A function $f: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ is compatible and preserves measure (respectively, is compatible and ergodic) iff it can be represented in the form $f(x) = c + x + 2v(x)$ (respectively, in the form $f(x) = 1 + x + 2\Delta v(x)$), where $c \in \mathbb{Z}_2$ and $v(x)$ is a compatible function.*

Proof. Follows easily from 2.1–2.3 and 2.5 in combination with (1) of the proof of the theorem 2.4. $\square$

Both 2.5–2.6 and theorem 2.7 could be applied to construct measure-preserving or ergodic functions as compositions of given compatible functions. For instance, putting $v(x) = (x^2) \text{ XOR } (x + 32 \text{ AND } x)$ (this function is compatible as a composition of compatible functions) we conclude that the function

$$7 + x + 2((x^2 + 2x + 1) \text{ XOR } (x + 1 + 32 \text{ AND } (x + 1))) - 2(x^2 \text{ XOR } (x + 32 \text{ AND } x))$$

is ergodic. This conclusion is not very easy to verify by direct application of theorems 2.2 or 2.3.

By the way, for $p = 2$ the statement of theorem 2.7 could be slightly modified to make it a little bit more convenient for the construction of ergodic functions out of addition and bitwise logical operations (like bitwise exclusive ‘or’, XOR, bitwise ‘and’, AND, or bitwise negation NEG). Namely, it could be easily seen that in the ring $\mathbb{Z}_2$ there holds an identity $z + \text{NEG}(z) = -1$. Hence, $\Delta v(x) = v(x+1) - v(x) = v(x+1) + \text{NEG}(v(x)) + 1$, and we obtain the following

2.8 Proposition. *A function $f: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ is compatible and ergodic iff it can be represented in one (hence, all) of the following forms $f(x) = 1 + x + 2(v(x+1) + \text{NEG}v(x)) = 2 + x + 2v(x+1) + \text{NEG}(2v(x)) = 3 + x + 2v(x+1) + 2\text{NEG}v(x)$, where $v: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ is an arbitrary compatible function.*

Since multiplication by 2 is just a 1-digit shift of 2-base expansion of a number towards senior bits, the proposition 2.8 could be applied to construct pseudorandom number generators out of the ‘fast’ computer commands, like addition, bitwise logical operations and shifts towards senior bits, by implementing the function v as a composition of them.

It worth noticing also that all the functions described in 2.4 – 2.8 are ‘affine modulo p ’, i.e., induce on $\mathbb{Z}/p$ a transformation of the form $x \mapsto a + bx$.

3. HENSEL LIFT.

This section studies conditions when a function of an important class of uniformly differentiable modulo p^k functions (which are properly defined below), is equiprobable, measure-preserving or ergodic. As a rule, the results of the section demonstrate

the effect of Hensel lift, mentioned in the introduction: speaking loosely, if a function F has some property modulo p^{k_0} then it has this property modulo p^n for all $n \geq k_0$. Besides, it worth noticing here that the results of this section, contrasting those of the previous one, provide some tools to construct measure-preserving or ergodic functions which are not necessarily affine modulo p . In fact, a certain techniques based on the ideas of this section could be developed; these techniques enables one ‘to lift’ an arbitrary transitive transformation of the ring $\mathbb{Z}/p^{k_0}$ to the function on $\mathbb{Z}_p$, which is transitive modulo p^k for all $k = k_0, k_0 + 1, k_0 + 2, \dots$. This is the main reason we introduce a notion of asymptotically compatible function below. However, the techniques themselves are not discussed here being left to the forthcoming paper.

Firstly, recall some generalizations of our basic notions (see 5.1 of [11]).

3.1 Definition. Let $F = (f_1, \dots, f_m): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$ be a function, not necessarily compatible. The function F is called (*asymptotically*) *equiprobable*, iff for all $k = 1, 2, \dots$ (respectively, for all sufficiently large $k \in \mathbb{N}$) it is *equiprobable modulo p^k* , that is, the restriction $F \bmod p^k = (f_1 \bmod p^k, \dots, f_m \bmod p^k)$ of the function F to the set $\{0, 1, \dots, p^k - 1\}^{(n)}$ is an equiprobable function. (Note that in cases which do not lead to misunderstanding we identify the set $\{0, 1, \dots, p^k - 1\}^{(n)}$ with the set of all elements of the ring $(\mathbb{Z}/p^k)^{(n)}$). By the analogy, we say that F is *asymptotically measure-preserving* (respectively, that F is *asymptotically ergodic*), iff $F \bmod p^k$ is a bijective (respectively, transitive) transformation of the ring $(\mathbb{Z}/p^k)^{(n)}$ for all sufficiently large k . Lastly, we say that F is *asymptotically compatible* iff there exists positive integer rational N such that for all $\mathbf{a}, \mathbf{b} \in \mathbb{Z}_p^{(n)}$ and all $k \geq N$ a congruence $\mathbf{a} \equiv \mathbf{b} \pmod{p^k}$ implies a congruence $F(\mathbf{a}) \equiv F(\mathbf{b}) \pmod{p^k}$.

By definition, for $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ of $\mathbb{Q}_p^{(n)}$ the congruence $\mathbf{a} \equiv \mathbf{b} \pmod{p^s}$ means that $\|a_i - b_i\|_p \leq p^{-s}$ (or, the same, that $a_i = b_i + c_i p^s$ for suitable $c_i \in \mathbb{Z}_p$, $i = 1, 2, \dots, s$); that is $\|\mathbf{a} - \mathbf{b}\|_p \leq p^{-s}$. In other words, a function is asymptotically compatible iff for some $N \in \mathbb{N}_0$ it satisfy Lipschitz condition with coefficient 1 for each pair of points which are at least as close one to another as p^{-N} . Since $\mathbb{Z}_p^{(n)}$ is compact, F is asymptotically compatible iff it satisfy Lipschitz condition with coefficient 1 locally.

Now for reader’s convenience we recall some facts of [11]. A function $F = (f_1, \dots, f_m): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$ is called *differentiable modulo p^k* at the point $\mathbf{u} = (u_1, \dots, u_n) \in \mathbb{Z}_p^{(n)}$, iff there exist a positive integer rational N and $n \times m$ matrix $F'_k(\mathbf{u})$ over $\mathbb{Q}_p$ (called a *Jacobi matrix modulo p^k* of the function F at the point $\mathbf{u}$) such that for each positive integer rational $K \geq N$ and each $\mathbf{h} = (h_1, \dots, h_n) \in \mathbb{Z}_p^{(n)}$ the inequality $\|\mathbf{h}\|_p \leq p^{-K}$ implies a congruence

$$F(\mathbf{u} + \mathbf{h}) \equiv F(\mathbf{u}) + \mathbf{h}F'_k(\mathbf{u}) \pmod{p^{k+K}}. \quad (\heartsuit)$$

In case $m = 1$ a Jacobi matrix modulo p^k is called a *differential modulo p^k* . In case $m = n$ a determinant of Jacobi matrix modulo p^k is called a *Jacobian modulo p^k* . The elements of Jacobi matrix modulo p^k are called *partial derivatives modulo p^k* of the function F at the point $\mathbf{u}$. A partial derivative (respectively, a differential) modulo p^k are sometimes denoted as $\frac{\partial_k f_i(\mathbf{u})}{\partial_k x_j}$ (respectively, as $d_k F(\mathbf{u}) = \sum_{i=1}^n \frac{\partial_k F(\mathbf{u})}{\partial_k x_i} d_k x_i$).

The definition immediately implies that partial derivatives modulo p^k of the function F are defined up to the p -adic integer summand which p -adic norm does not exceed p^{-k} . In cases when all partial derivatives modulo p^k at all points of $\mathbb{Z}_p^{(n)}$ are p -adic integers, we say that the function F has *integer-valued derivative modulo p^k* ; in these cases we can associate to each partial derivative modulo p^k a unique element of the ring $\mathbb{Z}/p^k$, and a Jacobi matrix modulo p^k at each point $\mathbf{u} \in \mathbb{Z}_p^{(n)}$ thus can be considered as a matrix over a ring $\mathbb{Z}/p^k$.

Under the latter agreement the ‘rules of differentiation modulo p^k ’ have the same (up to congruence modulo p^k instead of equality) form as for usual differentiation. For instance, if both functions $G: \mathbb{Z}_p^{(s)} \rightarrow \mathbb{Z}_p^{(n)}$ and $F: \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$ are differentiable modulo p^k at the points, respectively, $\mathbf{v} = (v_1, \dots, v_s)$ and $\mathbf{u} = G(\mathbf{v})$, and their partial derivatives modulo p^k at these points are p -adic integers, then a composition $F \circ G: \mathbb{Z}_p^{(s)} \rightarrow \mathbb{Z}_p^{(m)}$ of these functions is uniformly differentiable modulo p^k at the point $\mathbf{v}$, all its partial derivatives modulo p^k at this point are p -adic integers, and $(F \circ G)'_k(\mathbf{v}) \equiv G'_k(\mathbf{v})F'_k(\mathbf{u}) \pmod{p^k}$.

By the analogy with classical case we define for the function F a notion of *uniform differentiability modulo p^k on $\mathbb{Z}_p^{(n)}$* ; the least $K \in \mathbb{N}$ such that $(\heartsuit)$ holds simultaneously for all $\mathbf{u} \in \mathbb{Z}_p^{(n)}$, whereas $\|h_i\|_p \leq p^{-K}$, $(i = 1, 2, \dots, n)$, is denoted via $N_k(F)$. The latter number plays an important role in further considerations.

We recall that accordingly to 2.12 of [11] all partial derivatives modulo p^k of the uniformly differentiable modulo p^k function F are periodic functions with period $p^{N_k(F)}$. This in particular implies that each partial derivative modulo p^k can be considered as a function defined on $\mathbb{Z}/p^{N_k(F)}$. Moreover, if $F = (f_1, \dots, f_m): \mathbb{N}_0^{(n)} \rightarrow \mathbb{N}_0^{(m)}$ could be continued to a function on the $\mathbb{Z}_p^{(n)}$, which is uniformly differentiable modulo p^k on the whole space $\mathbb{Z}_p^{(n)}$, this continuation could be done simultaneously with all its (partial) derivatives modulo p^k .

Here and after in this section let $F = (f_1, \dots, f_m): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$ and $f: \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p$ be functions, which are uniformly differentiable on $\mathbb{Z}_p^{(n)}$ modulo p . This is relatively weak restriction since all uniformly differentiable on $\mathbb{Z}_p^{(n)}$ functions, as well as functions, which are uniformly differentiable on $\mathbb{Z}_p^{(n)}$ modulo p^k for some $k \geq 1$, are uniformly differentiable on $\mathbb{Z}_p^{(n)}$ modulo p .

The examples of functions which are not uniformly differentiable on $\mathbb{Z}_p^{(n)}$, yet are uniformly differentiable on $\mathbb{Z}_p^{(n)}$ modulo p , are the function $f(x, y) = x \text{ XOR } y$ for $p = 2$ and its corresponding analogs for $p \neq 2$; all partial derivatives modulo p of these functions are congruent to 1 modulo p at all points (see [11]). Note by the way, that previously introduced function $\text{mod } p^n: \mathbb{Z}_p \rightarrow \mathbb{Z}/p^n$, the ‘reduction modulo p^n ’, is uniformly differentiable on $\mathbb{Z}_p$ (its derivative is 0 at all points); the function $f(x, y) = x \text{ AND } y$ is differentiable modulo 2 at no point of $\mathbb{Z}_2^{(2)}$, yet it is uniformly differentiable with respect to x for each $y \in \mathbb{Z}$: its derivative is 0 for $y \geq 0$, and it is 1 in the opposite case.

It turns out that properties of being asymptotically compatible or asymptotically measure-preserving impose certain restrictions on p -adic norms of derivatives modulo p of a given function.

3.2 Proposition. *If the function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ asymptotically preserves measure, then $\|f'_1(u)\|_p \geq 1$ at all points $u \in \mathbb{Z}_p$.*

Proof. Since a derivative modulo p^k of the function f is periodic with period $p^{N_k(f)}$, it is sufficient to prove the proposition assuming $u \in \mathbb{N}_0$. Definition of differentiability modulo p^k implies that for $K \geq N_1(f)$ and for $u \in \mathbb{N}_0$ the congruence

$$f(u+h) \equiv f(u) + hf'_1(u) \pmod{p^{K+1}} \quad (1)$$

holds as soon as $\|h\|_p \leq p^{-K}$. Assuming $\|f'_1(u)\|_p < 1$ for some $u \in \mathbb{N}_0$, the condition $f'_1(u) \equiv 0 \pmod{p}$ and congruence (1) imply that $f(u+p^K) \equiv f(u) \pmod{p^{K+1}}$. The latter congruence means that for all $K \geq N_1(f)$, such that $u + p^K \leq p^{K+1} - 1$, the function f is not bijective modulo p^{K+1} . A contradiction. $\square$

3.3 Corollary. *If under the assumptions of 3.2 the function f is uniformly differentiable, then $\|f'(u)\|_p \geq 1$ for all $u \in \mathbb{Z}_p$.*

Proof. Definition of a derivative modulo p immediately implies that

$$f'_1(u) \equiv f'(u) \pmod{p}$$

for all $u \in \mathbb{Z}_p$. Thus $f'(u) = f'_1(u) + ps(u)$ for a suitable function $s: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$. Yet if $\|f'_1(u)\|_p \geq 1$, then the latter equality obviously implies that $\|f'(u)\|_p \geq 1$ by the properties of p -adic distance. Now the conclusion follows from 3.2. $\square$

The inverse of 3.2 is not true: an obvious counterexample gives the function $\frac{x^2-x}{2}$ on $\mathbb{Z}_2$. It vanishes both at 0 and at 1, but the 2-adic norm of its derivative is 2 everywhere on $\mathbb{Z}_2$. Nevertheless, functions of this kind are locally injective. Namely, the following is true:

3.4 Proposition. *If the function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is uniformly differentiable modulo p , and if $\|f'_1(u)\|_p \geq 1$, then a space $\mathbb{Z}_p$ can be represented as a disjoint union of a finite number of open (and simultaneously closed) balls U , for which the following holds: if $a, b \in U$, $k \geq N_1(f)$ and $a \not\equiv b \pmod{p^k}$, then $f(a) \not\equiv f(b) \pmod{p^k}$.*

Proof. Consider a union

$$\mathbb{Z}_p = \bigcup_{a=0}^{p^N-1} (a + p^N \mathbb{Z}_p),$$

where $N = N_1(f)$. Each set $U = a + p^N \mathbb{Z}_p$ is an open (and at the same time closed) ball of radius p^{-N} (see [3]). Let $u, v \in U$, and let $u \neq v$. Then $v = u + h$, where $\|h\|_p = p^{-K}$ for a suitable positive integer rational $K \geq N$. The definition of differentiability modulo p implies that

$$f(u+h) \equiv f(u) + hf'_1(u) \pmod{p^{K+1}}. \quad (1)$$

Thus, if $f(u) \equiv f(v) \pmod{p^K}$, then (1) implies that $\|f'_1(u)\|_p = p^{-1} < 1$. A contradiction. $\square$

The proposition 3.4 implies that if the p -adic norm of a uniformly differentiable modulo p function is not less than 1 everywhere on $\mathbb{Z}_p$, then this function might ‘glue together modulo p^k ’ for sufficiently large k only points which lie in distinct balls from the statement of 3.4. From here it follows

3.5 Proposition. *Let a function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be uniformly differentiable modulo p on $\mathbb{Z}_p$. Then f asymptotically preserves measure iff the following conditions hold simultaneously:*

- (1) $\|f'_1(u)\|_p \geq 1$ at all points $u \in \mathbb{Z}_p$;
- (2) $f(a) \not\equiv f(b) \pmod{p^n}$ for all $n, a, b \in \mathbb{N}_0$ such that $\|a - b\|_p \geq p^{-N_1(f)}$
 $0 \leq a, b \leq p^n - 1$. $\square$

A. A. Nechaev (private communication) noticed that the function $f(x) = \frac{x^2+x}{2}$ on $\mathbb{Z}_2$ asymptotically preserves measure (this also follows from 3.5). Thus, if a compatible function $g: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ asymptotically preserves measure (all these functions are characterized in 2.2), then a composition $h(x) = g(f(x))$ is uniformly differentiable modulo $p = 2$ and asymptotically measure-preserving function, and $\|g'_1(u)\|_2 = 2$ at all points $u \in \mathbb{Z}_2$. There are no other functions $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$, which are uniformly differentiable modulo p , asymptotically preserve measure, and which derivatives modulo p have norms not less than 1 everywhere on $\mathbb{Z}_p$, [10]. The proof of the latter statement involves not only p -adic tools, but algebraic geometry techniques as well.

The latter notice illustrates the fact that the second condition of the criterion 3.5 is rather difficult to verify since one has to calculate values of a function at infinite number of points. However, the problem might be simplified by imposing certain restrictions on the function under study. Namely, we will assume additionally that f maps each ball of radius p^{-M} (with $M \geq N_1(f)$) into a ball of radius p^{-M} (consequently, f is asymptotically compatible). This restriction is equivalent to the property of derivative modulo p to be integer-valued everywhere on $\mathbb{Z}_p$.

3.6 Proposition. *If for some $M \geq N_1(f)$ a uniformly differentiable modulo p function f maps each ball of radius p^{-M} into a ball of radius p^{-M} , then $f'_1(a) \in \mathbb{Z}_p$ for all $a \in \mathbb{Z}_p$. Vice versa, each uniformly differentiable modulo p function, which has an integer-valued derivative modulo p everywhere on $\mathbb{Z}_p$, maps each ball of radius p^{-M} into a ball of radius p^{-M} for all $M \geq N_1(f)$.*

Proof. If $M \geq N_1(f)$ and $\|h\|_p \leq p^{-M}$, then the definition of uniform differentiability modulo p^k (see 2.4 of [11]) implies that

$$f(u+h) \equiv f(u) + hf'_1(u) \pmod{p^{M+1}} \quad (1)$$

for all $u \in \mathbb{Z}_p$. On the other hand, the inclusion $f(a + p^M \mathbb{Z}_p) \subseteq f(a) + p^M \mathbb{Z}_p$ implies that

$$\|f(u+h) - f(u)\|_p \leq p^{-M} \quad (2)$$

for all h with $\|h\|_p \leq p^{-M}$. Comparing (1) and (2) we see that $\|f'_1(u)\|_p \leq 1$. The inverse statement is equivalent to the asymptotic compatibility of f (see 2.10 of [11]). $\square$

Henceforth in the section we additionally assume that f and F have integer-valued derivatives modulo p . In particular, this implies that both f and F are asymptotically compatible (see 2.10 and 2.11 of [11]). Now we state necessary and sufficient conditions the function F must satisfy to be measure-preserving, and sufficient conditions for equiprobability of F .

3.7 Theorem. *Let a function $F = (f_1, \dots, f_m): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(m)}$ be uniformly differentiable modulo p and let all its partial derivatives modulo p be integer-valued on $\mathbb{Z}_p$. Then F is asymptotically equiprobable if it is equiprobable modulo p^k for some $k \geq N_1(F)$ and the rank of its Jacobi matrix $F'_1(u)$ modulo p is exactly m at all points $\mathbf{u} = (u_1, \dots, u_n) \in (\mathbb{Z}/p^k)^{(n)}$.*

Proof. For $\xi \in (\mathbb{Z}/p^s)^{(m)}$ denote $F_s^{-1}(\xi) = \{\gamma \in (\mathbb{Z}/p^s)^{(n)} : F(\gamma) \equiv \xi \pmod{p^s}\}$. Let $s \geq k \geq N_1(F)$. Since F is asymptotically compatible, and hence F is a sum of a compatible function and a periodic function with period $p^{N_1(F)}$ (see 2.10 of [11]), we conclude that if $\eta \in F_{s+1}^{-1}(\xi)$, then $\bar{\eta} \in F_s^{-1}(\bar{\xi})$. Here, in accordance with our agreement in the introduction, $\bar{\alpha} = (\bar{\alpha}_1, \dots, \bar{\alpha}_m) \in (\mathbb{Z}/p^s)^{(m)}$ stands for $\alpha \bmod p^s = (\alpha_1 \bmod p^s, \dots, \alpha_m \bmod p^s)$, where $\alpha = (\alpha_1, \dots, \alpha_m) \in (\mathbb{Z}/p^{s+1})^{(m)}$. Put $\lambda = \bar{\eta} + p^s \sigma \in (\mathbb{Z}/p^{s+1})^{(n)}$, where $\sigma \in (\mathbb{Z}/p)^{(n)}$. In view of the uniform differentiability of the function F modulo p (see ($\heartsuit$)), we have

$$F(\lambda) \equiv F(\eta) + p^s \sigma F'_1(\bar{\eta}) \pmod{p^{s+1}}. \quad (1)$$

Since $F(\bar{\eta}) \equiv \bar{\xi} + p^k \beta \pmod{p^{s+1}}$ and $\xi = \bar{\xi} + p^s \gamma$ for suitable $\beta, \gamma \in (\mathbb{Z}/p)^{(m)}$, then (1) implies that $\lambda \in F_{s+1}^{-1}(\xi)$ iff $\bar{\lambda} \in F_s^{-1}(\bar{\xi})$ (i.e., $\bar{\eta} \in F_s^{-1}(\bar{\xi})$) and α satisfies the following linear system over a field $\mathbb{Z}/p$:

$$\beta + \alpha F'_1(\bar{\eta}) = \gamma. \quad (2)$$

Thus, if columns of the matrix $F'_1(\bar{\eta})$ are linearly independent over $\mathbb{Z}/p$, then linear system (2) has exactly p^{n-m} distinct solutions for arbitrary $\beta, \gamma \in (\mathbb{Z}/p)^{(m)}$. From here it follows that

$$|F_{s+1}^{-1}(\xi)| = |F_s^{-1}(\bar{\xi})| p^{n-m}. \quad (3)$$

Hence, if F is equiprobable modulo p^k (i.e., if $|F_s^{-1}(\bar{\xi})|$ does not depend on $\bar{\xi}$) and a rank of a matrix $F'_1(\bar{\eta})$ is m , then (3) implies that F is equiprobable modulo p^{s+1} . $\square$

3.8 Corollaries. 1° *Under the assumptions of theorem 3.7 let $m = 1$. Then F is asymptotically equiprobable if F is equiprobable modulo p^k for some $k \geq N_1(F)$, and differential $d_1 F$ modulo p of the function F vanishes at no point of $(\mathbb{Z}/p^k)^{(n)}$.*

2° *Let $f(x_1, \dots, x_n)$ be a polynomial with integer p -adic coefficients and in variables $x_1, \dots, x_n$. A polynomial f is equiprobable if it is equiprobable modulo p and all its partial derivatives vanishes simultaneously modulo p at no point of $(\mathbb{Z}/p)^{(n)}$ (i.e., are simultaneously congruent modulo p nowhere).*

Proof. The assertion 1° trivially follows from 3.7. In turn, 2° immediately follows from 1°, since for all $f \in \mathbb{Z}[x_1, \dots, x_n]$ holds $N_1(f) \leq 1$. We have only to prove the latter inequality.

By Taylor formula,

$$f(x_1 + h_1, \dots, x_n + h_n) = f(x_1, \dots, x_n) + \sum_{i=1}^n h_i \frac{\partial f}{\partial x_i} + Q \quad (1)$$

where $Q \in \mathbb{Z}[x_1, \dots, x_n, h_1, \dots, h_n]$, and each monomial in a canonic representation of the polynomial Q is of degree not less than 2 with respect to variables $h_1, \dots, h_n$. Since $\|(h_1, \dots, h_n)\|_p = p^{-s}$, where $s \geq 1$, for all values of $x_1, \dots, x_n$ we have $Q \equiv 0 \pmod{p^{2s}}$. In view of (1) this proves the inequality. $\square$

For $m = n$ the above stated sufficient conditions of asymptotical equiprobability occur to be necessary as well.

3.9 Theorem. Uniformly differentiable modulo p function

$$F = (f_1, \dots, f_m): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(n)}$$

with integer-valued derivatives modulo p , asymptotically preserves measure if and only if it is bijective modulo $p^{N_1(F)}$ and its Jacobian modulo p vanishes at no point of $(\mathbb{Z}/p^{N_1(F)})^{(n)}$ (equivalent condition: iff F is bijective modulo $p^{N_1(F)+1}$).

Proof. If F is bijective modulo $p^{N_1(F)}$ and its Jacobian modulo p vanishes nowhere, then in view of 3.7 F is asymptotically equiprobable, hence, asymptotically preserves measure, since $m = n$.

Vise versa, let F asymptotically preserves measure, i.e., let F be bijective modulo p^k for all $k \geq N$, where N is some positive integer rational. Now take $k \geq \max\{N, N_1(F)\}$, then the definition of uniform differentiability modulo p implies that

$$F(u + p^k \alpha) \equiv F(u) + p^k \alpha F'_1(u) \pmod{p^{k+1}} \quad (1)$$

for all $u, \alpha \in \mathbb{Z}_p$. Here $F'_1(u)$ is an $n \times n$ matrix over a field $\mathbb{Z}/p$. If $\det F'_1(u) \equiv 0 \pmod{p}$ for some $u \in \mathbb{Z}_p^{(n)}$ (or, the same, for some $u \in \{0, 1, \dots, p^{N_1(F)} - 1\}^{(n)}$ in view of the periodicity of partial derivatives modulo p), then there exists $\alpha \in \{0, 1, \dots, p-1\}^{(n)}$, $\alpha \not\equiv (0, \dots, 0) \pmod{p}$, such that $\alpha F'_1(u) \equiv (0, \dots, 0) \pmod{p}$. But then (1) implies that $F(u + p^k \alpha) \equiv F(u) \pmod{p^{k+1}}$. The latter contradicts the bijectivity modulo p^{k+1} of the function F , since for $u \in \{0, 1, \dots, p^{N_1(F)} - 1\}^{(n)}$ we have $u, u + p^k \alpha \in \{0, 1, \dots, p^{k+1} - 1\}^{(n)}$ and $u + p^k \alpha \neq u$.

Now we prove the criterion in the equivalent form. Let F be bijective modulo $p^{N_1(F)}$. Then assuming $k = N_1(F)$ in the above argument, we conclude that $\det F'_1(u) \not\equiv 0 \pmod{p}$ for all $u \in \mathbb{Z}_p^{(n)}$. According to 3.7, this implies that F asymptotically preserves measure.

Let F asymptotically preserves measure, and let it be not bijective modulo p^k for some $k \geq N_1(F)$. We prove that in this case F is not bijective modulo p^{k+1} .

Choose $u, v \in \{0, 1, \dots, p^k - 1\}^{(n)}$ such that $u \neq v$ $F(u) \equiv F(v) \pmod{p^k}$. Then either $F(u) \equiv F(v) \pmod{p^{k+1}}$ (i.e., F is not bijective modulo p^{k+1}), or $F(u) \not\equiv F(v) \pmod{p^{k+1}}$. Yet in the latter case we have $F(u) \equiv F(v) + p^k \alpha \pmod{p^{k+1}}$ for some $\alpha \in \{0, 1, \dots, p-1\}^{(n)}$, $\alpha \not\equiv (0, \dots, 0) \pmod{p}$. Consider $u_1 = u + p^k \beta$, where $\beta \in \{0, 1, \dots, p-1\}^{(n)}$ with $\beta \not\equiv (0, \dots, 0) \pmod{p}$ and $\beta F'_1(u) + \alpha \equiv (0, \dots, 0) \pmod{p}$. Such β exists, since F asymptotically preserves measure and, consequently, $\det F'_1(u) \not\equiv 0 \pmod{p}$, as it have been proven already. Now the definition of uniform differentiability modulo p implies that

$$F(u + p^k \beta) \equiv F(u) + p^k \beta F'_1(u) \equiv F(v) + p^k \alpha + p^k \beta F'_1(u) \equiv F(v) \pmod{p^{k+1}}, \quad (2)$$

where $u + p^k \beta \in \{0, 1, \dots, p^{k+1} - 1\}^{(n)}$ and $u + p^k \alpha \neq v$ (since $u \neq v$). Thus (2) in combination with our assumption imply that F is not bijective modulo p^{k+1} . Applying this argument sufficient number of times, we conclude that F is not bijective modulo p^s for all $s \geq k$. But at the same time F asymptotically preserves measure. A contradiction. $\square$

3.10 Corollaries. 1° If $n = 1$ within assumptions of the theorem 3.9, then F asymptotically preserves measure iff it is bijective modulo $p^{N_1(F)}$ and its derivative modulo p vanishes at no point of $\{0, 1, \dots, p^{N_1(F)} - 1\}$.

2° (cf. [8, Ch. 4, sections 4–5]) Let $F = (f_1, \dots, f_m): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(n)}$, where $f_i(x_1, \dots, x_n) \in \mathbb{Z}_p[x_1, \dots, x_n]$, $i = 1, 2, \dots, m$. Then F preserves measure iff F is bijective modulo p and $\det F'(u) \not\equiv 0 \pmod{p}$ for all $u \in \{0, 1, \dots, p-1\}^{(n)}$ (equivalent statement: iff F is bijective modulo p^2).

3° Let $A = \langle \mathbb{Z}_p; \Omega \rangle$ be a universal algebra of finite signature Ω , and let all operations of Ω are uniformly differentiable modulo p and have integer-valued derivatives modulo p . Then a polynomial over A defines an asymptotically measure-preserving function iff it is bijective modulo $p^{k(A)}$, where $k(A) = \max\{N_1(\omega) : \omega \in \Omega\} + 1$.

Proof. Assertion 1° trivially follows from 3.9. Assertion 2° holds in view of 3.9, since $N_1(F) \leq 1$ (see proof of the corollary 3.8). A composition $F \circ G$ of functions F and G , which are both uniformly differentiable modulo p and have integer-valued derivatives modulo p , is uniformly differentiable modulo p function, which has an integer-valued derivative modulo p , and $N_1(F \circ G) \leq \max\{N_1(F), N_1(G)\}$. The latter proves assertion 3°. $\square$

Comparing statements 3.7 and 3.9 one may ask a natural question whether sufficient conditions of 3.7 are necessary. The answer is negative: the results of [9] make it possible to construct the following counterexample.

Consider a function $f(x, y) = 2x + y^3$ on $\mathbb{Z}_2$. As f is a polynomial over $\mathbb{Z}$, then it is uniformly differentiable, has integer-valued derivatives, and $df = 2dx + 3y^2dy$. So, $df \equiv 0 \pmod{2}$ if $y \equiv 0 \pmod{2}$. Nevertheless, f induces an equiprobable function $(\mathbb{Z}/2^n)^{(2)} \rightarrow \mathbb{Z}/2^n$ for every $n = 1, 2, \dots$. Here is a proof.

For $n = 1$ we have that $f(x, y) \equiv y \pmod{2}$ is an equiprobable function on $\mathbb{Z}/2$. Let $n > 1$. We will show that for every $z \in \mathbb{Z}/2^n$ there exist exactly 2^n pairs (x, y) , such that $f(x, y) \equiv z \pmod{2^n}$ and $(x, y) \in \{0, 1, \dots, 2^n - 1\}^{(2)}$.

In fact, if $z = 1 + 2r$ for some $r \in \{0, 1, \dots, 2^{n-1} - 1\}$, then it follows that $y = 1 + 2k$ for some $k \in \{0, 1, \dots, 2^{n-1} - 1\}$. So $2x + (1 + 2k)^3 \equiv 1 + 2r \pmod{2^n}$ implies $x + 3k + 6k^2 + 4k^3 \equiv r \pmod{2^{n-1}}$. The left hand part of the latter congruence is a polynomial $\phi(x, k)$ in x, k . It is equiprobable in view of 3.8, 2°, since $d\phi \equiv dx + dk \pmod{2}$ (and hence this differential vanishes modulo 2 nowhere) and $\phi \equiv x + k \pmod{2}$ is obviously an equiprobable modulo 2 function. This implies that the congruence $\phi(x, k) \equiv r \pmod{2^{n-1}}$ in unknowns x, k has exactly 2^{n-1} solutions in $\{0, 1, \dots, 2^{n-1} - 1\}^{(2)}$.

If $z = 2r$ for some $r \in \{0, 1, \dots, 2^{n-1} - 1\}$, then it follows that $y = 2k$ for some $k \in \{0, 1, \dots, 2^{n-1} - 1\}$; consequently, the congruence $f(x, y) \equiv z \pmod{2^n}$ implies the congruence $x + 4k^3 \equiv r \pmod{2^{n-1}}$. Again the function $\psi(x, k)$ in the left hand part of the latter congruence is equiprobable in view of 3.8, 2°, since $d\psi \equiv dx \pmod{2}$ vanishes modulo 2 at no point of $(\mathbb{Z}/2)^{(2)}$ and $\psi \equiv x \pmod{2}$ is equiprobable modulo 2. From here, using an argument similar to one of the previous case, we conclude that the congruence $f(x, y) \equiv 2r \pmod{2^n}$ in unknowns x, y has exactly 2^n solutions in $\{0, 1, \dots, 2^n - 1\}^{(2)}$. Thus, f is equiprobable.

Now we are to begin a study of asymptotically ergodic functions in the class of all uniformly differentiable modulo p functions, which have integer-valued derivatives modulo p . It turns out that these functions could be in one variable only. To be more exact, the following theorem is true.

3.11 Theorem. Let a function $F = (f_1, \dots, f_n): \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(n)}$ be uniformly differentiable modulo p and asymptotically ergodic, and let it have integer-valued derivatives modulo p . Then $n = 1$.

We will need two lemmata.

3.12 Lemma. *Let a function $f: \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p$ be uniformly differentiable modulo p , let it have integer-valued derivatives modulo p , and let it vanish modulo p^k (i.e., let it be congruent 0 modulo p^k) for some $k > N_1(f)$ at all points of $\mathbb{Z}_p^{(n)}$. Then each partial derivative modulo p of the function f vanishes modulo p at all points of $\mathbb{Z}_p^{(n)}$.*

Proof of the lemma 3.12. Each function $g_i(x_0, x_1, \dots, x_n) = x_i + x_0 f(x_1, \dots, x_n)$ for arbitrary values of $x_0, x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n$ is a bijective modulo p^k function in variable x_i , ($i = 1, 2, \dots, n$). As $k > N_1(g_i) = N_1(f)$, then according to 3.9, g_i asymptotically preserves measure, and thus its derivative modulo p vanishes at no point of $\mathbb{Z}_p$. Moreover, the following is true:

$$\frac{\partial_1}{\partial_1 x_i} g_i(u_0, \dots, u_n) = 1 + u_0 \cdot \frac{\partial_1}{\partial_1 x_i} f(u_1, \dots, u_n) \not\equiv 0 \pmod{p} \quad (1)$$

for all $u_0, \dots, u_n \in \mathbb{Z}_p$. If

$$\frac{\partial_1}{\partial_1 x_i} f(u_1, \dots, u_n) \equiv d \not\equiv 0 \pmod{p}$$

for some $u_1, \dots, u_n \in \mathbb{Z}_p$, then choosing u_0 such that $u_0 d \equiv -1 \pmod{p}$ we get a contradiction to (1). This proves the lemma. $\square$

3.13 Lemma. *Let a function $H: \mathbb{Z}_p^{(n)} \rightarrow \mathbb{Z}_p^{(n)}$ be uniformly differentiable modulo p , and let it have integer-valued derivatives modulo p . If H is bijective modulo p^k and if H induces a trivial permutation modulo p^{k-1} (i.e., an identity transformation of $(\mathbb{Z}/p^{k-1})^{(n)}$) for some $k > N_1(H) + 1$, then H induces modulo p^k (i.e., on $(\mathbb{Z}/p^k)^{(n)}$) either a trivial permutation, or a permutation of order p .*

Proof of the lemma 3.13. Let G be an arbitrary function which satisfies assumptions of the lemma, and let $N_1(G) = N_1(H)$. Represent both H and G in the following form:

$$\begin{aligned} H(x_1, \dots, x_n) &= (x_1, \dots, x_n) + U(x_1, \dots, x_n); \\ G(x_1, \dots, x_n) &= (x_1, \dots, x_n) + V(x_1, \dots, x_n). \end{aligned}$$

Then both U and V are uniformly differentiable modulo p , have integer-valued derivatives modulo p , and $N_1(U) = N_1(V) = N_1(H)$. Moreover, both U and V vanish modulo p^{k-1} on $\mathbb{Z}_p^{(n)}$, for $k-1 > N_1(H)$. Then lemma 3.12 implies that $U'_1 = V'_1 = 0$ at all points of $\mathbb{Z}_p^{(n)}$. As $\|U\|_p \leq p^{-k+1}$ and $\|V\|_p \leq p^{-k+1}$ everywhere on $\mathbb{Z}_p^{(n)}$, then, implying 2.4, for all $h_1, \dots, h_n \in \mathbb{Z}_p$ we obtain, consequently, that

$$\begin{aligned} H(G(h_1, \dots, h_n)) &= H((h_1, \dots, h_n) + V(h_1, \dots, h_n)) \\ &\equiv H(h_1, \dots, h_n) + V(h_1, \dots, h_n) H'_1(h_1, \dots, h_n) \\ &\equiv H(h_1, \dots, h_n) + V(h_1, \dots, h_n) + V(h_1, \dots, h_n) U'_1(h_1, \dots, h_n) \\ &\equiv (h_1, \dots, h_n) + U(h_1, \dots, h_n) + V(h_1, \dots, h_n) \pmod{p^k}. \end{aligned}$$

This implies, in particular, that for all $s \in \mathbb{N}$ the following congruence holds:

$$\begin{aligned} H^s(h_1, \dots, h_n) &= \underbrace{H(\dots H(h_1, \dots, h_n) \dots)}_{s \text{ times}} \\ &\equiv (h_1, \dots, h_n) + sU(h_1, \dots, h_n) \pmod{p^k}. \end{aligned}$$

As U vanishes modulo p^{k-1} everywhere, then the latter congruence implies that $H^p(h_1, \dots, h_n) \equiv (h_1, \dots, h_n) \pmod{p^k}$ for all $h_1, \dots, h_n \in \mathbb{Z}_p$. This proves the lemma. $\square$

Proof of the theorem 3.11. Choose $k > N_1(F) + 1$ such that F is transitive modulo p^n for all $n \geq k - 1$. The function F induces a permutation on $(\mathbb{Z}/p^k)^{(n)}$ which is denoted as $\sigma_k(F)$. Consider a permutation $\sigma = \sigma_k(F)^{p^{(k-1)n}}$. As F is transitive modulo p^k , the order of σ is p^n (and hence σ is not trivial).

On the other hand, $\sigma = \sigma_k(F^{p^{(k-1)n}})$. But $F^{p^{(k-1)n}}$ is bijective modulo p^k and induces a trivial permutation modulo p^{k-1} (the latter assertion follows from transitivity of F modulo p^{k-1}). Since σ is not trivial, in view of 3.13 the order of σ must be p . Yet, according to the previous argument, the order of σ is p^n , so necessarily $n = 1$. $\square$

It is still an open problem to characterize asymptotically ergodic functions in the class of all uniformly differentiable modulo p functions which have integer-valued derivatives modulo p , but if we additionally assume that the function is uniformly differentiable modulo p^2 and has integer-valued derivative modulo p^2 , the following description can be obtained. The method we prove the next theorem is in fact a generalization to p -adic case of the idea originally applied by M. V. Larin to description of transitive modulo n polynomials over $\mathbb{Z}$, [15].

3.14 Theorem. *Let a function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be uniformly differentiable modulo p^2 and let it have integer-valued derivative modulo p^2 . Then f is asymptotically ergodic if and only if it is transitive modulo $p^{N_2(f)+1}$ for odd prime p or, respectively, modulo $2^{N_2(f)+2}$ for $p = 2$.*

We need the following

3.15 Lemma. *Let a function $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be uniformly differentiable modulo p , and let it have integer-valued derivative modulo p . If f is transitive modulo p^k for some $k > N_1(f)$, then f induces on $\mathbb{Z}/p^{k+1}$ a permutation, which is either a single cycle of length p^{k+1} , or a product of p pairwise disjoint cycles of length p^k each.*

Proof of the lemma 3.15. For $i = 0, 1, 2, \dots$ we denote via $x_i = \delta_i(x) \in \{0, 1, \dots, p-1\}$ a value of i th digit in canonic representation of p -adic integer $x \in \mathbb{Z}_p$. Now the definition of uniform differentiability modulo p implies that for an arbitrary $x \in \mathbb{Z}_p$ and $s \geq N_1(f) = N$ there holds a congruence $f(x_0 + x_1p + \dots + x_{s-1}p^{s-1} + x_sp^s) \equiv f(x_0 + x_1p + \dots + x_{s-1}p^{s-1}) + x_sp^s f'_1(x_0 + x_1p + \dots + x_{s-1}p^{s-1}) \pmod{p^{s+1}}$. The latter implies that

$$\delta_s(f(x)) \equiv \Phi_s(x_0, \dots, x_{s-1}) + x_s f'_1(x) \pmod{p}, \quad (1)$$

where $x_i = \delta_i(x) \in \{0, 1, \dots, p-1\}$ is the i -th p -adic digit of $x \in \mathbb{Z}_p$, ($i = 0, 1, 2, \dots$); $\Phi_s(x_0, \dots, x_{s-1}) = \delta_s(f(x_0 + x_1p + \dots + x_{s-1}p^{s-1}))$.

Since partial derivative $f'_1(x)$ modulo p is periodic with period p^N , it depends only on $x_0, \dots, x_{N-1}$, so (1) can be represented in the form

$$\delta_s(f(x)) \equiv \Phi_s(x_0, \dots, x_{s-1}) + x_s \Psi(x_0, \dots, x_{N-1}) \pmod{p}, \quad (2)$$

where $\Psi(x_0, \dots, x_{N-1}) = f'_1(x)$. Applying for the composition of functions ‘rules of differentiation modulo p^k ’ which were mentioned at the beginning of the section,

we conclude that for all $r = 1, 2, \dots$ the following congruence holds:

$$(f^r(x))'_1 \equiv \prod_{j=0}^{r-1} f'_1(f^j(x)) \pmod{p}. \quad (3)$$

We recall that $f^r(x) = \underbrace{f(\dots f(x) \dots)}_{r \text{ times}}$, $f^0(x) = x$. As f is asymptotically compatible, then transitivity of f modulo p^k for some $k \geq N$ implies transitivity of f modulo p^n for all $k \geq n \geq N$ (see [11], theorems 2.10 and 1.4). Yet f'_1 depends only on $x_0, \dots, x_{N-1}$, and f is transitive modulo p^N , so (3) implies that

$$(f^{p^n}(x))'_1 \equiv \left(\prod_{u_0, \dots, u_{N-1}=0}^{p-1} \Psi(u_0, \dots, u_{N-1}) \right)^{p^{n-N}} \pmod{p}. \quad (4)$$

We denote the product in the brackets in the right hand part of (4) as Π . Now, since $f^{p^n}(x)$ is uniformly differentiable modulo p and has integer-valued derivative modulo p , in view of (2) and (4) we conclude that

$$\delta_n(f^{p^n}(x)) \equiv \phi_n(x_0, \dots, x_{n-1}) + x_n \Pi^{p^{n-N}} \pmod{p}, \quad (5)$$

where $\phi_n(x_0, \dots, x_{n-1}) = \delta_n(f^{p^n}(x_0 + x_1 p + \dots + x_{n-1} p^{n-1}))$. Since f is a transitive modulo p^{n+1} function for $k \geq n \geq N$, the function f^{p^n} , on the one hand, induces a trivial permutation modulo p^n , and on the other hand, induces on each coset $a + p^n(\mathbb{Z}/p^{n+1})$ of the ring $\mathbb{Z}/p^{n+1}$ a permutation, which is a cycle of length p . This, in particular, means that the function in the right hand part of (5), being considered as a function in variable x_n , must be a permutation, moreover – a cycle of length p on $\{0, 1, \dots, p-1\}$. It is well known, however, that a polynomial $c + dy \in \mathbb{Z}[y]$ is transitive modulo p iff $d \equiv 1 \pmod{p}$ and $c \not\equiv 0 \pmod{p}$ (see e.g. [2, Ch. 3, Theorem A]). This implies, in particular, that $\Pi^{p^{n-N}} \equiv 1 \pmod{p}$, and hence $\Pi \equiv 1 \pmod{p}$. Finally we obtain that

$$\begin{aligned} f^{p^k}(x) &\equiv f^{p^k}(x_0 + x_1 p + \dots + x_k p^k) \\ &\equiv x_0 + x_1 p + \dots + x_{k-1} p^{k-1} + p^k(\phi_k(x_0, \dots, x_{k-1}) + x_k) \pmod{p^{k+1}}. \end{aligned} \quad (6)$$

The latter congruence implies that f induces a permutation σ modulo p^{k+1} . Moreover, we assert that if

$$\phi_k(x_0, \dots, x_{k-1}) \not\equiv 0 \pmod{p}$$

for some (equivalently, all) $x_0, \dots, x_{k-1} \in \{0, 1, \dots, p-1\}$, then f is transitive modulo p^{k+1} ; otherwise the permutation σ is a product of exactly p disjoint cycles of length p^k each.

To prove this assertion, consider some $u_0, \dots, u_k \in \{0, 1, \dots, p-1\}$ and denote C a cycle of the permutation σ which contains the point $u_0 + u_1 p + \dots + u_{k-1} p^{k-1} + x_k p^k \in \mathbb{Z}/p^{k+1}$. As f is transitive modulo p^k then (see (6)) p^k is a factor of $|C|$, the length of the cycle C . If $\phi_k(u_0, \dots, u_{k-1}) \not\equiv 0 \pmod{p}$, then (6) implies that

$$\begin{aligned} f^{p^k}(u_0 + u_1 p + \dots + u_{k-1} p^{k-1} + x_k p^k) \\ \not\equiv u_0 + u_1 p + \dots + u_{k-1} p^{k-1} + x_k p^k \pmod{p^{k+1}}, \end{aligned} \quad (7)$$

i.e., that $|C| > p^k$. On the other hand, (6) implies that $|C|$ is a factor of p^{k+1} . Finally we conclude that in this case $|C| = p^{k+1}$, i.e., f is transitive modulo p^{k+1} .

If $\phi_k(u_0, \dots, u_{k-1}) \equiv 0 \pmod{p}$ holds for some $u_0, \dots, u_k \in \{0, 1, \dots, p-1\}$, then this congruence holds for all $u_0, \dots, u_k \in \{0, 1, \dots, p-1\}$ (otherwise in view of the previous case f is transitive modulo p^{k+1} and (7) holds for all $u_0, \dots, u_k \in \{0, 1, \dots, p-1\}$ and the latter in view of (6) means that $\phi_k(u_0, \dots, u_{k-1}) \not\equiv 0 \pmod{p}$, a contradiction). Then (6) implies that σ^{p^k} is an identity permutation, i.e. $|C| = p^k$, as p^k is a factor of $|C|$. This proves the lemma. $\square$

Proof of the theorem 3.14. During the proof of the previous lemma we have established that if f is transitive modulo p^k for some $k \geq N_1(f)$, then f is transitive modulo p^n for all $k \geq n \geq N_1(f)$. So the ‘only if’ part of the theorem is proved, as $N_2(f) + 1 > N_1(f)$.

Now we have to prove that if $n \geq N_2(f) + 1$ (resp., if $n \geq N_2(f) + 2$ for $p = 2$) and if f is transitive modulo p^n , then it is transitive modulo p^{n+1} . In view of lemma 3.15 it is sufficient to prove that for some $x \in \mathbb{Z}_p$ the following condition holds:

$$f^{p^n}(x) \not\equiv x \pmod{p^{n+1}}. \quad (1)$$

As transitivity modulo p^n implies transitivity modulo p^{n-1} , in view of lemma 3.15 we have

$$f^{p^{n-1}}(x) = x + p^{n-1}\xi(x), \quad (2)$$

where $\xi: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ and $\xi(x) \not\equiv 0 \pmod{p}$ for all $x \in \mathbb{Z}_p$ (otherwise 3.15 implies that f is not transitive modulo p^n , a contradiction to the assumption).

Further, since f is uniformly differentiable modulo p^2 and has integer-valued derivative modulo p^2 , then for all $r = 1, 2, \dots$ a composition f^r is uniformly differentiable modulo p^2 and has integer-valued derivative modulo p^2 , and $(f^r(x))'_2 \equiv \prod_{j=0}^{r-1} f'_2(f^j(x)) \pmod{p^2}$ (see (3) of 3.15). Now, as $n-1 \geq N_2(f)$, then taking into account these considerations and an obvious (following from (2)) equality $f^{sp^{n-1}}(x) = f^{(s-1)p^{n-1}}(x + p^{n-1}\xi(x))$, where $s = 1, 2, \dots$, we successively calculate

$$\begin{aligned} f^{p^n}(x) &\equiv f^{(p-1)p^{n-1}}(x) + p^{n-1}\xi(x) \prod_{j=0}^{(p-1)p^{n-1}-1} f'_2(f^j(x)) \\ &\equiv f^{(p-2)p^{n-1}}(x) + p^{n-1}\xi(x) \left(\prod_{j=0}^{(p-2)p^{n-1}-1} f'_2(f^j(x)) + \prod_{j=0}^{(p-1)p^{n-1}-1} f'_2(f^j(x)) \right) \\ &\equiv \dots \equiv x + p^{n-1}\xi(x) \left(1 + \sum_{i=1}^{p-1} \prod_{j=0}^{(p-i)p^{n-1}-1} f'_2(f^j(x)) \right) \pmod{p^{n+1}}. \quad (3) \end{aligned}$$

Yet f'_2 is a periodic function with period $p^{N_2(f)}$ and f is transitive modulo p^{n-1} , so we conclude that for arbitrary $i, j \in \mathbb{N}$ the following congruence holds:

$$f'_2(f^j(x)) \equiv f'_2(f^{j+ip^{n-1}}(x)) \pmod{p^2}.$$

In view of the transitivity of f modulo p^{n-1} the latter congruence implies that

$$\prod_{j=0}^{(p-i)p^{n-1}-1} f'_2(f^j(x)) \equiv \alpha(x)^{p-i} \pmod{p^2},$$

where

$$\alpha(x) = \prod_{j=0}^{p^{n-1}-1} f'_2(f^j(x)).$$

In view of (3) we now conclude that

$$f^{p^n}(x) \equiv x + p^{n-1}\xi(x) \left(1 + \sum_{i=1}^{p-1} \alpha(x)^i \right) \pmod{p^{n+1}}. \quad (4)$$

Again, as f'_2 modulo p^2 is periodic with period $p^{N_2(f)}$ and f is transitive modulo p^{n-1} for $n-1 \geq N_2(f)$, then $\alpha(x)$ modulo p^2 does not depend on x . Moreover, we assert that $\alpha(x) \equiv 1 \pmod{p}$.

In fact, during the proof of 3.15 we have already established that if $k \geq N_1(f)$ and if f is a transitive modulo p^k and uniformly differentiable modulo p function with integer-valued derivative modulo p , then

$$\prod_{j=0}^{p^{N_1(f)}-1} f'_1(f^j(x)) \equiv 1 \pmod{p} \quad (5)$$

for all $x \in \mathbb{Z}_p$ (see the proof of (6) in 3.15). The definition of a derivative modulo p^2 implies that $f'_2(x) \equiv f'_1(x) \pmod{p}$; consequently,

$$\alpha(x) \equiv 1 + p\beta \pmod{p^2} \quad (6)$$

for some $\beta \in \mathbb{N}_0$. In view of (5) and (6), now (4) implies that

$$f^{p^n}(x) \equiv x + p^{n-1}\xi(x) \left(p + p\beta \sum_{i=1}^{p-1} i \right) \pmod{p^{n+1}}, \quad (7)$$

and for $p \neq 2$ we conclude that

$$f^{p^n}(x) \equiv x + p^n\xi(x) \pmod{p^{n+1}}.$$

In view of 3.15 the latter proves the theorem for $p \neq 2$, since $\xi(x) \not\equiv 0 \pmod{p}$ (see the text which follows (2)).

For the case $p = 2$, the congruence (7) implies that

$$f^{2^n}(x) \equiv x + 2^n(1 + \beta) \pmod{2^{n+1}} \quad (8)$$

and to finish the proof it is sufficient to show that β is even.

For $n \geq N_2(f) + 2$ the transitivity of f modulo 2^n implies that f is transitive modulo $2^{N_2(f)+2}$, so in view of the definition of a derivative modulo p^2 we have that

$$f^{2^N}(x + 2^N\xi) \equiv f^{2^N}(x) + 2^N\xi \prod_{j=0}^{2^N-1} f'_2(f^j(x)) \pmod{2^{N+2}} \quad (9)$$

for $N = N_2(f)$, $\xi \in \mathbb{Z}_2$. As f is transitive modulo 2^{N+2} , then for arbitrary $x \in \{0, 1, \dots, 2^N - 1\}$ and with ξ running over $\{0, 1, 2, 3\}$ the mapping

$$\phi_x: \xi \mapsto \delta_N(f^{2^N}(x + 2^N \xi)) + 2\delta_{N+1}(f^{2^N}(x + 2^N \xi))$$

is a cycle of length 4 on $\mathbb{Z}/4$. In view of (6),

$$\prod_{j=0}^{2^N-1} f'_2(f^j(x)) \equiv 1 + 2\beta \pmod{4};$$

so (9) implies that

$$\phi_x(\xi) \equiv c(x) + \xi(1 + 2\beta) \pmod{4}, \quad (10)$$

where $c(x) = \delta_N(f^{2^N}(x)) + 2\delta_{N+1}(f^{2^N}(x))$. But for each x the mapping ϕ_x is transitive modulo 4, so (10) in view of the above mentioned transitivity criterion for polynomials of degree 1 (see [2, Ch. 3, Theorem A]) implies that $\beta \equiv 0 \pmod{2}$. $\square$

Note. The analog of the theorem 3.14 generally does not hold for a function which is uniformly differentiable modulo p . Namely, for each $n \in \mathbb{N}$ there exists a uniformly differentiable modulo 2 and compatible function $f: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$ with $f'_1 = 1$ everywhere on $\mathbb{Z}_2$, $N_1(f) = 1$, which is transitive modulo 2^k for $k = 1, 2, \dots, n$, but which is not transitive modulo 2^k for all $k > n$. (By argument similar to applied below one can construct a counterexample for $p \neq 2$ as well.)

Represent $x \in \mathbb{Z}_2$ in its canonic form $x = x_0 + x_1 \cdot 2 + x_2 \cdot 2^2 + \dots$, $x_0, x_1, x_2, \dots \in \{0, 1\}$. Consider a function

$$f(x) = \sum_{i=0}^{\infty} \phi_i(x_0, \dots, x_i) \cdot 2^i,$$

where each $\phi_i(x_0, \dots, x_i)$ is a Boolean polynomial, which is linear with respect to variable x_i . In other words, $\phi_i(x_0, \dots, x_i) = \psi_i(x_0, \dots, x_{i-1}) + x_i$ in the factor-ring $\mathbb{Z}/2[x_0, \dots, x_i]/(x_0^2 - x_0, \dots, x_i^2 - x_i)$ of the ring $\mathbb{Z}/2[x_0, \dots, x_i]$ of all polynomials in variables $x_0, \dots, x_i$ over $\mathbb{Z}/2$ with respect to the ideal, generated by $x_0^2 - x_0, \dots, x_i^2 - x_i$ (we assume $\psi_0 = 1$). It is not difficult to see that this function f is compatible (see 3.9 of [11]). Direct calculations show that for arbitrary $s \in \mathbb{N}$ and $h \in \mathbb{Z}_2$ there holds a congruence $f(x + 2^s h) \equiv f(x) + 2^s h \pmod{2^{s+1}}$, i.e., that the function f is uniformly differentiable modulo 2, and $f'_1 = 1$ everywhere on $\mathbb{Z}_2$, with $N_1(f) = 1$.

Further, in the theory of Boolean functions there are well known sufficient and necessary conditions for transitivity modulo 2^n of the function f of the considered kind: namely, it is transitive modulo 2^n iff $\phi_i(x_0, \dots, x_i) = \psi_i(x_0, \dots, x_{i-1}) + x_i$ for $i = 1, 2, \dots, n-1$, where each Boolean polynomial $\psi_i(x_0, \dots, x_{i-1})$ for $i = 1, 2, \dots, n-1$ is of odd weight (that is, the number of all Boolean vectors, satisfying it, is odd) and $\psi_0 = 1$. (This result, which is known as transitivity modulo 2^n criterion for triangle transformations, belongs to mathematical folklore, so it is difficult to refer the originating paper, yet a proof can be found in, e.g., [11], see 4.8 there).

Now choosing for a given $n \in \mathbb{N}$ a function f so that $\psi_0 = 1$, with Boolean polynomials $\psi_i(x_0, \dots, x_{i-1})$ of odd weight for $i = 1, 2, \dots, n-1$, and with Boolean polynomial $\psi_n(x_0, \dots, x_{n-1})$ of even weight, we obtain a function, which is transitive modulo 2^k for $k = 1, 2, \dots, n$, but which is not transitive modulo 2^{n+1} . Then it is not transitive each modulo 2^k with $k > n$, since, in view of compatibility of f , transitivity of f modulo 2^{k+1} implies its transitivity modulo 2^k .

3.16 Corollary. *Let $A = \langle \mathbb{Z}_p; \Omega \rangle$ be a universal algebra of finite signature Ω , and let all operations of Ω be uniformly differentiable modulo p^2 functions with integer-valued derivatives modulo p^2 . Then there exists a positive rational integer $k(A)$ such that a polynomial $f(x) \in A[x]$ is asymptotically ergodic if and only if it is transitive modulo $p^{k(A)}$.*

Proof. The proof of this corollary is similar to one of 3.10, 3° and so is omitted. We can take $k(A) = \max\{N_2(\omega) : \omega \in \Omega\} + \epsilon$, where $\epsilon = 1$ if p is odd, otherwise $\epsilon = 2$. $\square$

4. HENSEL LIFT STARTING POINTS.

The results of previous section show that for a class $\mathcal{D}_1$ (respectively, $\mathcal{D}_2$) of all uniformly differentiable modulo p (respectively, modulo p^2) functions, which have integer-valued derivatives modulo p (respectively, modulo p^2), there exists a function $\zeta: \mathcal{D}_1 \rightarrow \mathbb{N}$ (respectively, $\eta: \mathcal{D}_2 \rightarrow \mathbb{N}$), such that a function $f \in \mathcal{D}_1$ (respectively, $f \in \mathcal{D}_2$) is asymptotically measure-preserving (or is ergodic) iff it is bijective (respectively, transitive) modulo $p^{\zeta(f)}$ (respectively, modulo $p^{\eta(f)}$). Theorems 3.9 and 3.14 give corresponding estimates for $\zeta(f)$ and $\eta(f)$.

These estimates are sharp, i.e., there exist a compatible function $f \in \mathcal{D}_1$ (respectively, $f \in \mathcal{D}_2$) such that f is bijective (respectively, transitive) modulo $p^{N_1(f)}$ (respectively, modulo $p^{N_2(f)}$ for $p \neq 2$, or modulo $2^{N_2(f)+1}$ for $p = 2$), but f is not measure-preserving (respectively, is not ergodic). For instance, a polynomial $f(x) = 1 + x^p$ is bijective modulo p , $N_1(f) = 1$, but in force of 3.10, 1° the polynomial f is not bijective modulo p^2 , since $f'(z) \equiv 0 \pmod{p}$ for all $z \in \mathbb{Z}_p$.

A corresponding example for theorem 3.14 in case $p \neq 2$ gives a function $f(x) = (x+1) \odot_p 1$, where $\odot_p$ is digitwise multiplication modulo p of p -adic integers: $\delta_i(x \odot_p y) \equiv \delta_i(x) \delta_i(y) \pmod{p}$ for all $i \in \mathbb{N}_0$. The function f is uniformly differentiable, its derivative is 0 everywhere on $\mathbb{Z}_p$, and $N_2(f) = 1$; at the same time f is transitive modulo p , but it is not even bijective (hence, is not transitive) modulo p^2 .

Nevertheless, boundaries for $\zeta(f)$ and $\eta(f)$, which give, respectively, theorems 3.9 and 3.14, might differ significantly from the ones for various proper subclasses of $\mathcal{D}_1$ and of $\mathcal{D}_2$. For instance, for a function $f(x) = (ax + b) \text{ XOR } c$, with $a, b, c \in \mathbb{N}$, theorem 3.14 states that $f(x)$ is asymptotically ergodic iff it is transitive modulo $2^{\lfloor \log_2 c \rfloor + 2}$, since this function is uniformly differentiable and has a derivative which is a everywhere on $\mathbb{Z}_2$, and $N_2(f) = \lfloor \log_2 c \rfloor$. Yet direct application of the above mentioned criteria of transitivity modulo 2^n for triangle transformations and for polynomials of degree 1 over $\mathbb{Z}$ immediately implies that f is ergodic iff it is transitive modulo 4. So the problem of sharpening estimates of $\zeta(f)$ and $\eta(f)$ for various important from a certain view classes, which are narrower than $\mathcal{D}_1$ and $\mathcal{D}_2$, could be of interest.

In this section we study a class $\mathcal{A}$ of all compatible functions $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ such that, loosely speaking, coefficients of their interpolation series tends to 0 as fast as

$i!$, or faster (recall that $\lim_{i \rightarrow \infty}^p i! = 0$). More accurately, a function f , represented by interpolation series $(\diamond)$ (see section 2) with p -adic integer coefficients a_i , belongs to $\mathcal{A}$ iff it is compatible, and a sequence $\{\| \frac{a_i}{i!} \|_p : i = 0, 1, 2, \dots\}$ is bounded, i.e., $\| \frac{a_i}{i!} \|_p \leq p^{\rho(f)}$ for some $\rho(f) \in \mathbb{N}_0$. Recall that according to the theorem 2.1, a function f represented by $(\diamond)$ is compatible iff $\|a_i\|_p \leq p^{-\lfloor \log_p i \rfloor}$ for all $i \in \mathbb{N}$.

Class $\mathcal{A}$ is rather wide: it contains all integer-valued compatible analytic on $\mathbb{Z}_p$ functions, in particular, compatible functions which could be defined by integer-valued polynomials over $\mathbb{Q}_p$. It is known (see [3, Ch. 4, Theorem 4, p. 224]), that a function f of the form $(\diamond)$ is analytic on $\mathbb{Z}_p$ iff $\lim_{i \rightarrow \infty}^p \frac{a_i}{i!} = 0$.

So for the rest of this section we assume that $f \in \mathcal{A}$. Put

$$\lambda(f) = \min \left\{ k \in \mathbb{N} : 2 \frac{p^k - 1}{p - 1} - k > \rho(f) \right\}.$$

The following theorem is true.

4.1 Theorem. *Let $f \in \mathcal{A}$ and p is an odd prime. The function f is ergodic if and only if it is transitive modulo $p^{\lambda(f)+1}$ (if $p \neq 3$) or modulo $3^{\lambda(f)+2}$ (if $p = 3$).*

Since f is compatible, then in view of 2.1 it can be represented in the following form:

$$f(x) = b_0 + \sum_{i=1}^{\infty} b_i p^{\lfloor \log_p i \rfloor} \binom{x}{i},$$

where $b_j \in \mathbb{Z}_p$ for $j = 0, 1, 2, \dots$. Everywhere during the proof we assume that f is represented in this form. Further $\lambda(f)$ is denoted as λ and p is assumed to be an odd prime. We will need some additional technical results.

4.2 Lemma. *Under the assumptions of theorem 4.1 the following is true:*

$$\begin{aligned} b_i &\equiv 0 \pmod{p}, \text{ for } i \geq 2p^\lambda; \\ b_i &\equiv 0 \pmod{p^2}, \text{ for } i \geq 3p^\lambda. \end{aligned}$$

Proof of the lemma 4.2. If $b_i = 0$, then the assertion of the lemma is trivial. Suppose that $b_i \neq 0$. Represent f as

$$f(x) = b_0 + \sum_{i=1}^{\infty} \frac{1}{i!} b_i p^{\lfloor \log_p i \rfloor} (x)_i,$$

where, we recall, $(x)_i = x(x-1) \cdots (x-i+1)$ (with $(x)_0 = 1$) is i th descending factorial power of x . As $f \in \mathcal{A}$, i.e.,

$$\left\| b_i p^{\lfloor \log_p i \rfloor} \right\|_p \leq p^{\rho(f)} \|i!\|_p$$

then

$$\text{ord}_p b_i \geq \text{ord}_p i! - \lfloor \log_p i \rfloor - \rho(f), \quad (1)$$

for all $i = 1, 2, \dots$. We recall that $\log_p \|a\|_p = -\text{ord}_p a$, for $a \in \mathbb{Z}_p$. Thus, the maximal p -prime factor of a is exactly $p^{\text{ord}_p a}$.

In fact, the function $\kappa(i) = \text{ord}_p i! - \lfloor \log_p i \rfloor$ is nondecreasing. To prove this, note that, obviously, $\text{ord}_p i! \geq \text{ord}_p (i-1)!$. If $\lfloor \log_p i \rfloor = \lfloor \log_p (i-1) \rfloor$ then $\kappa(i-1) \leq \kappa(i)$.

Assume $\lfloor \log_p j \rfloor > \lfloor \log_p (j-1) \rfloor$ for some positive rational integer j . Evidently, $\lfloor \log_p j \rfloor + 1$ is the number of significant digits in the p -base expansion of j . Hence the case under consideration takes place exactly if and only if $j-1 = (p-1) + (p-1)p + \dots + (p-1)p^n = p^{n+1} - 1$ for some $n \in \mathbb{N}_0$. But then $\text{ord}_p j! = \text{ord}_p (j-1)! + n$, $\lfloor \log_p (j-1) \rfloor = n$, $\lfloor \log_p j \rfloor = n+1$, and so $\kappa(j) > \kappa(j-1)$.

Now it is sufficient to prove only that $\kappa(2p^\lambda) - \rho(f) \geq 1$ and $\kappa(3p^\lambda) - \rho(f) \geq 2$. We recall that $\text{ord}_p i! = \frac{1}{p-1}(i - \text{wt}_p i)$, where $\text{wt}_p i$ is the sum of all digits in a p -base expansion of i (i.e., if $i = i_0 + i_1p + \dots + i_s p^s$, where $i_0, \dots, i_s \in \{0, 1, \dots, p-1\}$, then $\text{wt}_p i = i_0 + \dots + i_s$, see e.g., [6, ch.1, section 2, exercise 13]).

As $p \neq 2$, then $\kappa(2p^\lambda) - \rho(f) = \frac{1}{p-1}(2p^\lambda - 2) - \lambda - \rho(f) \geq 1$ according to the definition of $\lambda = \lambda(f)$. Hence, if $p \neq 3$, then

$$\kappa(3p^\lambda) - \rho(f) = \frac{1}{p-1}(3p^\lambda - 3) - \lambda - \rho(f) = \kappa(2p^\lambda) + \frac{1}{p-1}(p^\lambda - 1) - \rho(f) \geq 2.$$

So if $p \neq 3$ the lemma is proved.

Finally, let $p = 3$. Then

$$\kappa(3p^\lambda) - \rho(f) = \kappa(3^{\lambda+1}) - \rho(f) = \frac{1}{2}(3^{\lambda+1} - 1) - \lambda - 1 - \rho(f) \geq 2,$$

otherwise in view of the inequality

$$3^\lambda - 1 - \lambda > \rho(f),$$

(which follows directly from the definition of $\lambda = \lambda(f)$) we get

$$\frac{1}{2}(3^{\lambda+1} - 1) - \lambda - 1 - 3^\lambda + 1 + \lambda < 1,$$

i.e., $3^\lambda - 1 < 2$, and so $\lambda < 1$, a contradiction. The lemma 4.2 is proved. $\square$

4.3 Corollary. *Under assumptions of theorem 4.2, for $i \in \mathbb{N}$ the following is true:*

$$\frac{\Delta^i f(x)}{i} \equiv \begin{cases} 0 \pmod{p^2}, & \text{if } i \geq 2p^\lambda + 1; \\ 0 \pmod{p}, & \text{if } i \geq p^\lambda + 1. \end{cases}$$

Proof of the corollary 4.3. As $\Delta^j \binom{x}{i} = \binom{x}{i-j}$ if $i \geq j$ and $\Delta^j \binom{x}{i} = 0$ if $i < j$, then

$$\frac{\Delta^i f(x)}{i} = \frac{1}{\hat{i}} \sum_{j=i}^{\infty} b_j p^{\lfloor \log_p j \rfloor - \text{ord}_p j} \binom{x}{j-i},$$

where $\hat{i} = ip^{-\text{ord}_p i} \in \mathbb{Z}_p$, $\text{ord}_p \hat{i} = 0$. Now the result is obvious in view of lemma 4.2. $\square$

4.4 Proposition. *Under assumptions of theorem 4.1 the function f is uniformly differentiable modulo p^2 , has integer-valued derivative modulo p^2 , $N_2(f) \leq \lambda(f) + 1$. Moreover,*

$$f'_2(x) \equiv \sum_{i=1}^{2p^\lambda} (-1)^{i-1} \frac{\Delta^i f(x)}{i} \pmod{p^2}.$$

Proof of the proposition 4.4. To prove the first assertion of the proposition we will demonstrate that there exists a function $f'_2: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ such that for all $x, h \in \mathbb{Z}_p$ and $m \geq \lambda(f) + 1$ the following congruence holds:

$$f(x + p^m h) \equiv f(x) + p^m h f'_2(x) \pmod{p^{m+2}}. \quad (1)$$

In view of the compatibility of f , it is sufficient to prove the congruence (1) only for $h \in \{1, 2, \dots, p^2 - 1\}$ (for $h = 0$ the congruence is trivial). Applying Newton formula

$$f(x + n) = \sum_{i=0}^n \binom{n}{i} \Delta^i f(x)$$

for $n = p^m h$, we have

$$f(x + p^m h) = f(x) + p^m h \phi_m(x, h), \quad (2)$$

where

$$\phi_m(x, h) = \sum_{i=1}^{p^m h} \binom{p^m h - 1}{i - 1} \frac{\Delta^i f(x)}{i}. \quad (3)$$

Hence in view of 4.3 for $m \geq \lambda + 1$ we obtain:

$$\phi_m(x, h) \equiv \sum_{i=1}^{2p^\lambda} \binom{p^m h - 1}{i - 1} \frac{\Delta^i f(x)}{i} \pmod{p^2}. \quad (4)$$

Further, for $i = 1, 2, \dots, 2p^\lambda$ the following obvious equalities hold:

$$\binom{p^m h - 1}{i - 1} = \prod_{k=0}^{i-2} \frac{p^m h - (k + 1)}{k + 1} = \prod_{j=1}^{i-1} \left(\frac{h}{\hat{j}} p^{m - \text{ord}_p j} - 1 \right). \quad (5)$$

Here $\hat{j} = jp^{-\text{ord}_p j}$ is the unit of $\mathbb{Z}_p$, i.e., $\hat{j}$ has multiplicative inverse $\frac{1}{\hat{j}}$ in $\mathbb{Z}_p$; hence, each factor of the product in the right hand part of (5) is p -adic integer.

If $i \leq p^\lambda$ then $m - \text{ord}_p j \geq 2$ for all $j = 1, 2, \dots, i - 1$; so (5) implies that

$$\binom{p^m h - 1}{i - 1} \equiv (-1)^{i-1} \pmod{p^2}. \quad (6)$$

If $p^\lambda + 1 \leq i \leq 2p^\lambda$ and $j \in \{1, 2, \dots, i - 1\}$ then $m - \text{ord}_p j = 1$ only in the case when simultaneously $j = p^\lambda$ and $m = \lambda + 1$ hold; otherwise $m - \text{ord}_p j \geq 2$. Yet if $m - \text{ord}_p j = 1$ then

$$\frac{\Delta^i f(x)}{i} \equiv 0 \pmod{p}$$

(see 4.3); hence in both cases we have that

$$\left(\frac{h}{j}p^{m-\text{ord}_p j} - 1\right) \frac{\Delta^i f(x)}{i} \equiv -\frac{\Delta^i f(x)}{i} \pmod{p^2}.$$

So in view of (5) we conclude that

$$\left(\frac{p^m h - 1}{i - 1}\right) \frac{\Delta^i f(x)}{i} \equiv (-1)^{i-1} \frac{\Delta^i f(x)}{i} \pmod{p^2}. \quad (7)$$

for all $i = 1, 2, \dots, 2p^\lambda$. Now (4), (6), (7) together imply that

$$\phi_m(x, h) \equiv \sum_{i=1}^{2p^\lambda} (-1)^{i-1} \frac{\Delta^i f(x)}{i} \pmod{p^2}$$

and in view of (2), (3), (4) this completes the proof of proposition 4.4. $\square$

4.5. Lemma. *Under assumptions of theorem 4.1, there exists a function $\theta: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ such that for arbitrary $x, h \in \mathbb{Z}_p$ the following congruence holds:*

$$f(x + p^\lambda h) \equiv f(x) + p^\lambda h f'_2(x) + p^{\lambda+1} h^2 \theta(x) \pmod{p^{\lambda+2}}.$$

The function θ satisfies the following condition: for arbitrary $a, b \in \mathbb{Z}_p$ the congruence $a \equiv b \pmod{p^\lambda}$ implies $\theta(a) \equiv \theta(b) \pmod{p}$. Moreover, one may put

$$\theta(x) = \sum_{j=2}^{p-1} (-1)^j \sum_{i=1}^{j-1} \frac{1}{i} \frac{\Delta^{jp^{\lambda-1}} f(x)}{jp^{\lambda-1}} + \sum_{k=1}^{p-1} (-1)^{k-1} \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^\lambda} + \frac{\Delta^{2p^\lambda} f(x)}{2p^{\lambda+1}}.$$

Proof of the lemma 4.5. Firstly we prove that the function θ defined by the latter equality is integer-valued on $\mathbb{Z}_p$. Since f is compatible, each fraction $\frac{\Delta^s f(x)}{s}$ for $s = 1, 2, 3, \dots$, is p -adic integer (see 3.1 of [11]). So it is sufficient to prove only that for all $k \in \{1, 2, \dots, p-1\}$ both functions $\alpha(x)$ and $\beta_k(x)$ (defined below) are integer-valued on $\mathbb{Z}_p$. By definition,

$$\alpha(x) = \frac{\Delta^{2p^\lambda} f(x)}{2p^{\lambda+1}}; \quad \beta_k(x) = \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^\lambda}.$$

Since

$$\Delta^i f(x) = \sum_{j=i}^{\infty} b_j p^{\lfloor \log_p j \rfloor} \binom{x}{j-i} \quad (1)$$

for $i = 1, 2, 3, \dots$ and

$$b_j p^{\lfloor \log_p j \rfloor} \equiv 0 \pmod{p^{\lambda+1}}$$

for all integer rationals $j \geq 2p^\lambda$ (see 4.2), then $\alpha(x) \in \mathbb{Z}_p$. If $j \geq kp^{\lambda-1} + p^\lambda$ then $\lfloor \log_p j \rfloor \geq \lambda$; hence (1) implies that $\beta_k(x) \in \mathbb{Z}_p$.

Now we prove that for all $a, b \in \mathbb{Z}_p$ the congruence $a \equiv b \pmod{p^\lambda}$ implies $\theta(a) \equiv \theta(b) \pmod{p}$. In view of (1) and 4.2 the following congruence holds:

$$\alpha(x) \equiv \frac{1}{2} \sum_{j=2p^\lambda}^{3p^\lambda-1} \frac{1}{p} b_j \binom{x}{j-2p^\lambda} \pmod{p}. \quad (2)$$

We recall a statement of the well known Lucas theorem (for a proof see e.g [4]): if $a = \sum_{i=0}^{\infty} a_i p^i$ and $b = \sum_{i=0}^N b_i p^i$ are, respectively, canonic representations of p -adic integer a and of nonnegative integer rational b (i.e., $a_i, b_i \in \{0, 1, \dots, p-1\}$ for $i = 0, 1, 2, \dots$), then

$$\binom{a}{b} \equiv \binom{a_0}{b_0} \binom{a_1}{b_1} \cdots \binom{a_N}{b_N} \pmod{p}.$$

So, if $a \equiv b \pmod{p^\lambda}$, then Lucas theorem implies that for all $j = 2p^\lambda, 2p^\lambda + 1, \dots, 3p^\lambda - 1$ the following congruence holds:

$$\binom{a}{j-2p^\lambda} \equiv \binom{b}{j-2p^\lambda} \pmod{p}.$$

Thus, (2) implies that

$$\alpha(a) \equiv \alpha(b) \pmod{p}. \quad (3)$$

Further, combining (1) and 4.2 we obtain that

$$\beta_k(x) \equiv \frac{1}{k} \sum_{j=kp^{\lambda-1}+p^\lambda}^{2p^\lambda-1} b_j \binom{x}{j-kp^{\lambda-1}-p^\lambda} \pmod{p}$$

for all $k = 1, 2, \dots, p-1$. Now applying Lucas theorem once again, we conclude that

$$\beta_k(a) \equiv \beta_k(b) \pmod{p} \quad (4)$$

for $a \equiv b \pmod{p^\lambda}$.

Lastly, assuming

$$\gamma_k(x) = \frac{\Delta^{kp^{\lambda-1}} f(x)}{kp^{\lambda-1}},$$

in view of (1) we conclude that for $k = 1, 2, \dots, p-1$ the following congruence holds:

$$\gamma_k(x) \equiv \frac{1}{k} \sum_{j=kp^{\lambda-1}}^{p^\lambda-1} b_j \binom{x}{j-kp^{\lambda-1}} \pmod{p}.$$

Again, applying Lucas theorem, we conclude that

$$\gamma_k(a) \equiv \gamma_k(b) \pmod{p} \quad (5)$$

for $a \equiv b \pmod{p^\lambda}$. Hence in view of (3) – (5) the congruence $a \equiv b \pmod{p^\lambda}$ implies the congruence $\theta(a) \equiv \theta(b) \pmod{p}$.

Now we prove the rest of the lemma. As f is compatible, during the proof we may assume that $h \in \mathbb{N}$ (case $h = 0$ is trivial). According to 4.4 (see (2)–(5) there) the following is true:

$$f(x + p^\lambda h) \equiv f(x) + p^\lambda h \phi(x, h) \pmod{p^{\lambda+2}}, \quad (6)$$

where

$$\phi(x, h) \equiv \sum_{i=1}^{2p^\lambda} \binom{p^\lambda h - 1}{i-1} \frac{\Delta^i f(x)}{i} \pmod{p^2}. \quad (7)$$

and, besides,

$$\binom{p^\lambda h - 1}{i-1} = \prod_{j=1}^{i-1} \left(\frac{h}{j} p^{\lambda - \text{ord}_p j} - 1 \right) \quad (8)$$

for $i = 1, 2, \dots, 2p^\lambda$.

As f is compatible, then, according to 3.40 of [11],

$$\frac{\Delta^i f(x)}{i} \equiv 0 \pmod{p}$$

in all cases with the exception of, possibly, a case when i is of the form $i = tp^s$ for suitable $t \in \{1, 2, \dots, p-1\}$ and $s \in \mathbb{N}_0$. Thus, if $i \leq p^{\lambda-1}$, as well as if simultaneously $p^{\lambda-1} < i < p^\lambda$ and $p^{\lambda-1}$ is not a factor of i , the equality (8) implies:

$$\binom{p^\lambda h - 1}{i-1} \frac{\Delta^i f(x)}{i} \equiv (-1)^{i-1} \frac{\Delta^i f(x)}{i} \pmod{p^2}. \quad (9)$$

Let $i = kp^{\lambda-1}$ for $k \in \{2, 3, \dots, p-1\}$. Then (8) implies:

$$\binom{p^\lambda h - 1}{i-1} \equiv (-1)^{kp^{\lambda-1}-1} + (-1)^k p h \sum_{j=1}^{k-1} \frac{1}{j} \pmod{p^2}. \quad (10)$$

Further, if $p^\lambda \leq i \leq 2p^\lambda$ and $\text{ord}_p i \neq \lambda, \lambda-1$ then (1) (together with congruence following it) imply that

$$\frac{\Delta^i f(x)}{i} \equiv 0 \pmod{p^2}. \quad (11)$$

Now we have to study the only two remaining cases: $i = \nu p^\lambda$ for $\nu \in \{1, 2\}$ and $i = kp^{\lambda-1} + p^\lambda$ for $k \in \{1, 2, \dots, p-1\}$. The latter one in view of 4.3 and (8) implies that

$$\binom{p^\lambda h - 1}{i-1} \frac{\Delta^i f(x)}{i} \equiv (-1)^{i-1} \frac{\Delta^i f(x)}{i} + (-1)^{k-1} h \frac{\Delta^i f(x)}{i} \pmod{p^2}. \quad (12)$$

Further, for $k = 1, 2, \dots, p-1$ the following trivial equality holds in $\mathbb{Q}_p$:

$$\left(1 + \frac{p}{k}\right) \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1} + p^\lambda} = \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1}} \quad (13)$$

From here in view of 4.3 we conclude that

$$\frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1}+p^\lambda} \equiv 0 \pmod{p}$$

and since $\frac{p}{k} \in \mathbb{Z}_p$ and $\text{ord}_p \frac{p}{k} = 1$, the equality (13) implies that

$$\frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1}+p^\lambda} \equiv \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1}} \pmod{p^2}.$$

Hence, applying (12) for $i = kp^{\lambda-1} + p^\lambda$, we have that

$$\begin{aligned} & \left(\frac{p^\lambda h - 1}{kp^{\lambda-1} + p^\lambda - 1} \right) \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1} + p^\lambda} \\ &= (-1)^{kp^{\lambda-1}+p^\lambda-1} \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1} + p^\lambda} + (-1)^{k-1} ph\beta_k(x) \pmod{p^2}. \end{aligned} \quad (14)$$

In case $i = p^\lambda$, the equality (8) implies that

$$\left(\frac{p^\lambda h - 1}{p^\lambda - 1} \right) \equiv (-1)^{p^\lambda-1} - ph \sum_{j=1}^{p-1} \frac{1}{j} \equiv (-1)^{p^\lambda-1} \pmod{p^2}, \quad (15)$$

since for $p \neq 2$ the following congruences hold in $\mathbb{Q}_p$: $\sum_{j=1}^{p-1} \frac{1}{j} \equiv \sum_{j=1}^{p-1} j \equiv 0 \pmod{p}$.

Finally, for $i = 2p^\lambda$, applying (8) and 4.3, we conclude that

$$\begin{aligned} & \left(\frac{p^\lambda h - 1}{2p^\lambda - 1} \right) \frac{\Delta^{2p^\lambda} f(x)}{2p^\lambda} \equiv (-1)^{2p^\lambda-1} \frac{\Delta^{2p^\lambda} f(x)}{2p^\lambda} + h \frac{\Delta^{2p^\lambda} f(x)}{2p^\lambda} \\ & \equiv (-1)^{2p^\lambda-1} \frac{\Delta^{2p^\lambda} f(x)}{2p^\lambda} + hp\alpha(x) \pmod{p^2}, \end{aligned} \quad (16)$$

where $\alpha(x) \in \mathbb{Z}_p$, as it was shown above.

Now by the combination of (6), (7), (9), (11), (14), (15), (16) with 4.4 we finish the proof of the lemma 4.5. $\square$

4.6 Lemma. *Under assumptions of theorem 4.1, for all $x, h \in \mathbb{Z}_p$ the following congruence holds:*

$$f'_2(x + p^\lambda h) \equiv f'_2(x) + 2ph\theta(x) \pmod{p^2}.$$

Here θ is the function defined in 4.5.

Proof of the lemma 4.6. In view of 4.4 the following is true:

$$f'_2(x + p^\lambda h) \equiv \sum_{i=1}^{2p^\lambda} (-1)^{i-1} \frac{\Delta^i f(x + p^\lambda h)}{i} \pmod{p^2}. \quad (1)$$

For $i = 1, 2, \dots, 2p^\lambda$ the previous lemma implies that

$$\begin{aligned} \frac{\Delta^i f(x + p^\lambda h)}{i} &\equiv \frac{\Delta^i f(x)}{i} + hp^{\lambda - \text{ord}_p i} \frac{\Delta^i f'_2(x)}{\hat{i}} \\ &\quad + h^2 p^{\lambda + 1 - \text{ord}_p i} \frac{\Delta^i \theta(x)}{\hat{i}} \pmod{p^2}, \end{aligned} \quad (2)$$

where $\hat{i} = ip^{-\text{ord}_p i}$ is a unit in $\mathbb{Z}_p$, i.e., it has a multiplicative inverse $\frac{1}{\hat{i}} \in \mathbb{Z}_p$.

The term of order 2 (with respect to h) in (2) may not vanish modulo p^2 only if $i \in \{p^\lambda, 2p^\lambda\}$. Yet, as $\Delta^j \binom{x}{\nu} = \binom{x}{\nu-j}$ for $\nu \geq j$ and $\Delta^j \binom{x}{\nu} = 0$ for $\nu < j$, then for all $j \in \mathbb{N}$ we have

$$\Delta^j f(x) = \sum_{\nu=j}^{\infty} b_\nu p^{\lfloor \log_p \nu \rfloor} \binom{x}{\nu-j}. \quad (3)$$

Consequently, if $j \in \{p^\lambda, 2p^\lambda\}$, then

$$\frac{\Delta^{j+kp^{\lambda-1}} f(x)}{kp^{\lambda-1}} \equiv 0 \pmod{p}. \quad (4)$$

for $k \in \{1, 2, \dots, p-1\}$. Further, for $j \in \{p^\lambda, 2p^\lambda\}$ the equality (3) in view of 4.3 implies that

$$\frac{\Delta^{j+kp^{\lambda-1}+p^\lambda} f(x)}{kp^\lambda} \equiv 0 \pmod{p}, \quad (5)$$

$$\frac{\Delta^{j+2p^\lambda} f(x)}{2p^\lambda} \equiv 0 \pmod{p}. \quad (6)$$

Now, by the definition of θ , combining together (4), (5), (6) we conclude that $\frac{\Delta^i \theta(x)}{i} \equiv 0 \pmod{p}$ for $i \in \{p^\lambda, 2p^\lambda\}$, and thus

$$h^2 p^{\lambda+1 - \text{ord}_p i} \frac{\Delta^i \theta(x)}{\hat{i}} \equiv 0 \pmod{p^2} \quad (7)$$

for all $i = 1, 2, \dots, 2p^\lambda$.

The term of order 1 in (2) may not vanish modulo p^2 only for $i \in \{1, 2, \dots, 2p^\lambda\}$ such that $\text{ord}_p i \geq \lambda - 1$, i.e., for

$$i \in \{p^\lambda, 2p^\lambda, kp^{\lambda-1}, kp^{\lambda-1} + p^\lambda : k = 1, 2, \dots, p-1\}.$$

Combining together 4.3, 4.4 and 3.4 of [11] we already referred (see argument which follows (8) in the proof of 4.5), we have

$$f'_2(x) \equiv \frac{\Delta^{p^\lambda} f(x)}{p^\lambda} + \sum_{t=0}^{\lambda-1} \sum_{\tau=1}^{p-1} (-1)^{\tau-1} \frac{\Delta^{\tau p^t} f(x)}{\tau p^t} \pmod{p}, \quad (8)$$

and hence

$$\Delta^i f'_2(x) \equiv \frac{\Delta^{i+p^\lambda} f(x)}{p^\lambda} + \sum_{t=0}^{\lambda-1} \sum_{\tau=1}^{p-1} (-1)^{\tau-1} \frac{\Delta^{i+\tau p^t} f(x)}{\tau p^t} \pmod{p}. \quad (9)$$

This for $i \in \{kp^{\lambda-1} + p^\lambda : k = 1, 2, \dots, p-1\}$ in force of (3) and 4.2 implies that $\Delta^i f'_2(x) \equiv 0 \pmod{p}$, and consequently

$$hp \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f'_2(x)}{k+p} \equiv 0 \pmod{p^2} \quad (10)$$

for $k = 1, 2, \dots, p-1$ (since multiplicative inverse $\frac{1}{k+p}$ of $k+p$ is in $\mathbb{Z}_p$).

If $i \in \{kp^{\lambda-1} : k = 1, 2, \dots, p-1\}$ then in view of 4.2, (3) and (9) we have:

$$\Delta^{kp^{\lambda-1}} f'_2(x) \equiv \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{p^\lambda} + \sum_{\tau=1}^{p-k-1} (-1)^{\tau-1} \frac{\Delta^{(\tau+k)p^{\lambda-1}} f(x)}{\tau p^{\lambda-1}} \pmod{p}. \quad (11)$$

If $i = 2p^\lambda$ then 4.4 implies that

$$\Delta^{2p^\lambda} f'_2(x) \equiv \sum_{j=1}^{2p^\lambda} (-1)^{j-1} \frac{\Delta^{j+2p^\lambda} f(x)}{j} \pmod{p^2}.$$

This in view of (3) and 4.2 implies that

$$\Delta^{2p^\lambda} f'_2(x) \equiv 0 \pmod{p^2}. \quad (12)$$

Now we consider a case $i = p^\lambda$. Proposition 4.4 implies that

$$\Delta^{p^\lambda} f'_2(x) \equiv \sum_{j=1}^{1+p^\lambda} (-1)^{j-1} \frac{\Delta^{j+p^\lambda} f(x)}{j} \pmod{p^2}, \quad (13)$$

since, combining together (3) and 4.2, for $j = p^\lambda + 1, \dots, 2p^\lambda$ we conclude that

$$\frac{\Delta^{j+p^\lambda} f(x)}{j} \equiv 0 \pmod{p^2}.$$

Moreover, (3) implies that the latter congruence holds also for all $j \leq p^\lambda - 1$, such that $j \neq kp^{\lambda-1}$, where $k = 1, 2, \dots, p-1$. Thus, (13) implies that

$$\Delta^{p^\lambda} f'_2(x) \equiv \frac{\Delta^{2p^\lambda} f(x)}{p^\lambda} + \sum_{k=1}^{p-1} (-1)^{k-1} \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1}} \pmod{p^2}. \quad (14)$$

Now, substituting (7), (10), (11), (12), (14) to (2) and summarizing up all the obtained congruences for i ranging from 1 to $2p^\lambda$, in view of (1) and 4.4 we conclude that

$$\begin{aligned} f'_2(x + p^\lambda h) &\equiv f'_2(x) + hp \left(\sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \sum_{\tau=1}^{p-k-1} (-1)^{\tau-1} \frac{\Delta^{(\tau+k)p^{\lambda-1}} f(x)}{\tau p^{\lambda-1}} \right. \\ &\quad \left. + \sum_{k=1}^{p-1} (-1)^{k-1} \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^\lambda} \right) \\ &\quad + h \sum_{k=1}^{p-1} (-1)^{k-1} \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1}} + h \frac{\Delta^{2p^\lambda} f(x)}{p^\lambda} \pmod{p^2}. \end{aligned} \quad (15)$$

We recall that here and after all calculations are performed in the field $\mathbb{Q}_p$, and by the above agreement the congruence $\xi \equiv 0 \pmod{p^k}$ for $\xi \in \mathbb{Q}_p$ and positive integer rational k means that $\|\xi\|_p = p^{-k}$ (hence, ξ is a p -adic integer). Proceeding with this note, we conclude that for $k, \tau \in \{1, 2, \dots, p-1\}$ the following equalities hold in $\mathbb{Q}_p$:

$$\begin{aligned} & \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \sum_{\tau=1}^{p-k-1} (-1)^{\tau-1} \frac{\Delta^{(\tau+k)p^{\lambda-1}} f(x)}{\tau p^{\lambda-1}} \\ &= \sum_{m=1}^{p-1} (-1)^m \sum_{k+\tau=m} \frac{1}{k\tau} \cdot \frac{\Delta^{mp^{\lambda-1}} f(x)}{p^{\lambda-1}} = 2 \sum_{m=1}^{p-1} (-1)^m \sum_{\tau=1}^{m-1} \frac{1}{\tau} \cdot \frac{\Delta^{mp^{\lambda-1}} f(x)}{mp^{\lambda-1}}, \end{aligned} \quad (16)$$

since for $k, \tau \in \{1, 2, \dots, p-1\}$ it is obvious that

$$\sum_{k+\tau=m} \frac{1}{k\tau} = \sum_{k+\tau=m} \frac{1}{(m-\tau)\tau} = \frac{1}{m} \sum_{k+\tau=m} \left(\frac{1}{\tau} + \frac{1}{m-\tau} \right) = \frac{2}{m} \sum_{\tau=1}^{m-1} \frac{1}{\tau}.$$

Besides, as it was shown during the proof of 4.5, both $\alpha(x)$ and $\beta_k(x)$ are p -adic integers for $k = 1, 2, \dots, p-1$ and $x \in \mathbb{Z}_p$; thus

$$2hp\alpha(x) = h \frac{\Delta^{2p^\lambda} f(x)}{p^\lambda}; \quad hp\beta_k(x) = h \frac{\Delta^{kp^{\lambda-1}+p^\lambda} f(x)}{kp^{\lambda-1}}, \quad (17)$$

where all the factors are p -adic integers. Now the assertion of the lemma follows from (15), (16), (17) and definition of the function θ . $\square$

Proof of the theorem 4.1. Finishing the proof of the theorem 4.1, note that according to 4.4 there holds an inequality $N_2(f) \leq \lambda(f) + 1$. Thus, by 3.14 it is sufficient only to show that if $p \neq 3$ and f is transitive modulo $p^{\lambda(f)+1}$ then it is transitive modulo $p^{\lambda(f)+2}$. In turn, for this purpose in view of 3.15 it is sufficient only to prove that

$$f^{p^{\lambda+1}}(x) \not\equiv x \pmod{p^{\lambda+2}} \quad (1)$$

at least for one $x \in \mathbb{Z}_p$. Further we merely calculate $f^{p^{\lambda+1}}(x) \pmod{p^{\lambda+2}}$.

Under the assumptions we have made above, f is transitive modulo p^λ , since f is compatible. Then by 3.15 we conclude that for all $x \in \mathbb{Z}_p$

$$f^{p^\lambda}(x) = x + p^\lambda \xi(x), \quad \xi(x) \not\equiv 0 \pmod{p}, \quad (2)$$

where $\xi: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a function defined everywhere on $\mathbb{Z}_p$.

We assert that for each $i = 0, 1, 2, \dots$ the following congruence holds:

$$\begin{aligned} f^{p^\lambda+i}(x) &\equiv f^i(x) + p^\lambda \xi(x) \prod_{j=0}^{i-1} f'_2(f^j(x)) \\ &+ p^{\lambda+1} \xi(x)^2 \prod_{j=0}^{i-1} f'_2(f^j(x)) \sum_{k=0}^{i-1} \frac{\theta(f^k(x))}{f'_2(f^k(x))} \prod_{\tau=0}^{k-1} f'_2(f^\tau(x)) \pmod{p^{\lambda+2}} \end{aligned} \quad (3)$$

Recall that the sum (resp., product) over the empty set of indexes is assumed to be 0 (resp., 1). Note also that since f is transitive modulo $p^{\lambda+1}$ it is bijective modulo $p^{\lambda+1}$. Consequently, f is bijective modulo $p^\lambda, \dots, p^2, p$ since f is compatible. Hence $f'_1(x) \not\equiv 0 \pmod{p}$ for all $x \in \mathbb{Z}_p$ (see the proof of 3.9) and thus $f'_2(x) \not\equiv 0 \pmod{p}$ either (as $f'_2(x) \equiv f'_1(x) \pmod{p}$). So all the denominators in (3) have multiplicative inverses in $\mathbb{Z}_p$; thus, during the proof of (3) and further we assume that all the calculations are performed in $\mathbb{Z}_p$.

We can easily prove (3) by the induction on i . If $i = 0$, then (3) trivially follows from (2). Assume that (3) is true for $i = m - 1$. As

$$f^{p^\lambda+m}(x) = f(f^{p^\lambda+m-1}(x)) \quad (4)$$

then, assuming in (3) that $i = m - 1$, substituting (3) to (4), applying 4.5 and a congruence $(f^k(x))'_2 \equiv \prod_{j=0}^{k-1} f'_2(f^j(x)) \pmod{p^2}$, we prove the congruence (3) for $i = m$, in view of compatibility of f , by obvious direct calculations. We omit details.

Now we apply (3) to calculate $f^{p^{\lambda+1}}(x) \pmod{p^{\lambda+2}}$. Put

$$\begin{aligned} A_i(x) &= (f^i(x))'_2 = \prod_{j=0}^{i-1} f'_2(f^j(x)); \\ B_i(x) &= (f^i(x))'_2 \sum_{k=0}^{i-1} \frac{(f^k(x))'_2}{f'_2(f^k(x))} \theta(f^k(x)) = \\ &= \left(\prod_{j=0}^{i-1} f'_2(f^j(x)) \right) \cdot \left(\sum_{k=0}^{i-1} \frac{\theta(f^k(x))}{f'_2(f^k(x))^2} \prod_{\tau=0}^k f'_2(f^\tau(x)) \right). \end{aligned}$$

Lemma 4.6 implies that

$$f'_2(a + p^\lambda h) \equiv \begin{cases} f'_2(a) \pmod{p^2}, & \text{if } h = 0; \\ f'_2(a) \pmod{p}, & \text{if } h \neq 0. \end{cases} \quad (5)$$

As f is transitive modulo p^λ , then (5) implies that $f'_2(f^k(x)) \equiv f'_2(f^r(x)) \pmod{p}$ as soon as $k \equiv r \pmod{p^\lambda}$. Besides, by 4.5 the latter condition implies that $\theta(f^k(x)) \equiv \theta(f^r(x)) \pmod{p}$.

Further,

$$\prod_{\tau=0}^{p^\lambda-1} f'_2(f^\tau(x)) \equiv 1 \pmod{p}. \quad (6)$$

This has been already proven in 3.15 (see proof of (6) there), since 4.5 implies that $N_1(f) \leq \lambda$. Consequently,

$$\prod_{\tau=0}^k f'_2(f^\tau(x)) \equiv \prod_{\tau=0}^r f'_2(f^\tau(x)) \pmod{p}$$

as soon as $k \equiv r \pmod{p^\lambda}$.

Finally we conclude that for every $t \in \mathbb{N}$

$$B_{tp^\lambda}(x) \equiv t \sum_{\tau=0}^{p^\lambda-1} \frac{\theta(f^\tau(x))}{f_2'(f^\tau(x))^2} \prod_{\nu=0}^{\tau} f_2'(f^\nu(x)) \equiv tB_{p^\lambda}(x) \pmod{p}. \quad (7)$$

Now we calculate $A_{tp^\lambda}(x) \pmod{p^2}$ for $t \in \mathbb{N}$. The congruence (3) in view of (6) implies that

$$f^{kp^\lambda+\tau}(x) \equiv f^\tau(x) + kp^\lambda \xi(x) \prod_{j=0}^{\tau-1} f_2'(f^j(x)) \pmod{p^{\lambda+1}} \quad (8)$$

for all $k \in \mathbb{N}$ and all $\tau \in \{0, 1, \dots, p^\lambda - 1\}$. As

$$A_{tp^\lambda}(x) = \prod_{k=0}^{t-1} \prod_{\tau=0}^{p^\lambda-1} f_2'(f^{kp^\lambda+\tau}(x)),$$

then in view of (5) and 4.6 the congruence (8) implies that

$$A_{tp^\lambda}(x) = \prod_{k=0}^{t-1} \prod_{\tau=0}^{p^\lambda-1} f_2' \left(f^\tau(x) + kp^\lambda \xi(x) \prod_{j=0}^{\tau-1} f_2'(f^j(x)) \right) \pmod{p^2},$$

or, applying 4.6,

$$\begin{aligned} A_{tp^\lambda}(x) &= \prod_{k=0}^{t-1} \prod_{\tau=0}^{p^\lambda-1} \left(f_2'(f^\tau(x)) + 2kp\xi(x)\theta(f^\tau(x)) \prod_{j=0}^{\tau-1} f_2'(f^j(x)) \right) \\ &\equiv \prod_{k=0}^{t-1} \left(\prod_{\tau=0}^{p^\lambda-1} f_2'(f^\tau(x)) \right. \\ &\quad \left. + 2kp\xi(x) \sum_{s=0}^{p^\lambda-1} \theta(f^s(x)) \frac{\prod_{j=0}^{p^\lambda-1} f_2'(f^j(x))}{f_2'(f^s(x))} \prod_{j=0}^{s-1} f_2'(f^j(x)) \right) \pmod{p^2}. \end{aligned} \quad (9)$$

According to (6),

$$\prod_{j=0}^{p^\lambda-1} f_2'(f^j(x)) = 1 + p\epsilon$$

for suitable $\epsilon \in \mathbb{Z}_p$; consequently, (9) implies that

$$\begin{aligned} A_{tp^\lambda}(x) &\equiv \prod_{k=0}^{t-1} \left(1 + p\epsilon + 2kp\xi(x) \sum_{s=0}^{p^\lambda-1} \theta(f^s(x)) \frac{\prod_{j=0}^{s-1} f_2'(f^j(x))}{f_2'(f^s(x))} \right) \equiv \\ &\equiv 1 + tp\epsilon + 2p\xi(x) \left(\sum_{k=0}^{t-1} k \right) \cdot \left(\sum_{s=0}^{p^\lambda-1} \theta(f^s(x)) \frac{\prod_{j=0}^{s-1} f_2'(f^j(x))}{f_2'(f^s(x))^2} \right) \equiv \\ &\equiv 1 + tp\epsilon + pt(t-1)\xi(x) \sum_{s=0}^{p^\lambda-1} \theta(f^s(x)) \frac{\prod_{j=0}^{s-1} f_2'(f^j(x))}{f_2'(f^s(x))^2} \pmod{p^2}. \end{aligned} \quad (10)$$

Now combining together (2), (3), (7) and (10) we conclude that

$$\begin{aligned} f^{(t+1)p^\lambda}(x) &\equiv f^{tp^\lambda+p^\lambda}(x) \\ &\equiv f^{tp^\lambda}(x) + p^\lambda \xi(x) + \epsilon p^{\lambda+1} \xi(x) + p^{\lambda+1} t^2 \xi(x)^2 B_{p^\lambda}(x) \pmod{p^{\lambda+2}}. \end{aligned} \quad (11)$$

Finally, combining (11), (2) with obvious induction on n we obtain that

$$\begin{aligned} f^{np^\lambda}(x) &\equiv x + np^\lambda \xi(x) + \epsilon p^{\lambda+1} \xi(x) \frac{n(n-1)}{2} \\ &\quad + p^{\lambda+1} \xi(x)^2 B_{p^\lambda}(x) \frac{n(n-1)(2n-1)}{6} \pmod{p^{\lambda+2}} \end{aligned}$$

or, in particular,

$$f^{p^{\lambda+1}}(x) \equiv x + p^{\lambda+1} \xi(x) \pmod{p^{\lambda+2}},$$

since $p \neq 2, 3$. But the latter congruence in view of (2) implies that

$$f^{p^{\lambda+1}}(x) \not\equiv x \pmod{p^{\lambda+2}}.$$

This finally proves the theorem 4.1 $\square$

Note. With the use of theorem 4.1 we can determine whether a given integer-valued and compatible polynomial $f(x) \in \mathbb{Q}_p[x]$ is ergodic. Represent $f(x)$ in the form $f(x) = \frac{g(x)}{r}$, where $r \in \mathbb{Z}_p$ and $g(x) \in \mathbb{Z}_p[x]$ and at least one coefficient of $g(x)$ is coprime with p . In fact, we can take r to be a common denominator of all coefficients of $f(x)$ represented as irreducible fractions. Here we assume that $f(x)$ is represented in the basis $(x)_0 = 1, (x)_1 = x, (x)_2 = x(x-1), \dots$ of descending factorial powers, or in a standard basis $1, x, x^2, \dots$. Then $\rho(f) = \text{ord}_p r$, and $\rho(f)$ does not depend on the choice of the basis. We recall that $p^{\text{ord}_p r}$ is the greatest power of p which is a factor of r . Now we easily find $\lambda(f)$ and determine whether f is transitive on $\mathbb{Z}/p^{\lambda(f)+1}$ (e.g., by direct calculations). In view of 4.1 for $p \neq 2, 3$ this is equivalent to the ergodicity of $f(x)$ (for $p = 3$ one should study transitivity of f on $\mathbb{Z}/p^{\lambda(f)+2}$).

Moreover, it is possible for each prime p to determine, whether a polynomial $f(x) \in \mathbb{Q}_p[x]$ is integer-valued, compatible and ergodic, by calculating its values at $O(\deg f)$ points. Namely, the following is true.

4.7 Proposition. *A polynomial $f(x) \in \mathbb{Q}_p[x]$ is integer-valued, compatible and ergodic iff the mapping*

$$z \mapsto f(z) \pmod{p^{\lfloor \log_p(\deg f) \rfloor + 3}},$$

with z ranging over $\{0, 1, \dots, p^{\lfloor \log_p(\deg f) \rfloor + 3} - 1\}$, defines a compatible and transitive function on the residue class ring $\mathbb{Z}/p^{\lfloor \log_p(\deg f) \rfloor + 3}$.

Proof. Coefficients $a_i \in \mathbb{Q}_p$ ($i = 0, 1, \dots, d$) of the polynomial $f(x)$ of degree d , which is represented in the form $f(x) = \sum_{i=0}^d a_i \binom{x}{i}$ (see $(\diamond)$ of section 2), are defined by the values this polynomial $f(x)$ takes at the points $0, 1, \dots, d$. In other words, all values $f(0), f(1), \dots, f(d)$ are p -adic integers iff all coefficients $a_i \in \mathbb{Q}_p$ ($i = 0, 1, \dots, d$) are p -adic integers, i.e., iff a polynomial $f(x)$ is integer-valued (see

the beginning of section 2). By the analogy, in view of the theorem 2.1, a polynomial $f(x)$ preserves all congruences of the ring $\mathbb{Z}/p^{\lfloor \log_p d \rfloor + 1}$ iff $\|a_i\| \leq p^{-\lfloor \log_p i \rfloor}$ for all $i = 1, 2, \dots, d$, i.e., iff $f(x)$ is compatible on $\mathbb{Z}_p$. In other words, to determine whether a polynomial $f(x)$ is integer-valued and compatible it is sufficient (and necessary) to determine whether it induces a compatible function on the ring $\mathbb{Z}/p^k$ for some (arbitrarily fixed) $k \geq \lfloor \log_p d \rfloor + 1$.

In force of theorem 4.1, for $p \neq 2$, an integer-valued and compatible polynomial $f(x)$ is ergodic iff it is transitive modulo p^k for any arbitrarily fixed $k \geq \lambda(f) + 2$. Representing $f(x)$ as $f(x) = b_0 + \sum_{i=1}^d b_i p^{\lfloor \log_p i \rfloor} \binom{x}{i}$, $b_j \in \mathbb{Z}_p$ for $j = 0, 1, 2, \dots$, we conclude that $\rho(f)$ is the least nonnegative integer rational, which is not less than each of $\text{ord}_p i! - \lfloor \log_p i \rfloor - \text{ord}_p b_i$ ($i = 1, 2, \dots, d$). Thus, since a function $\text{ord}_p i! - \lfloor \log_p i \rfloor$ is nondecreasing (see proof of the lemma 4.2), then each $k \in \mathbb{N}$, which satisfies inequality $2 \frac{p^k - 1}{p - 1} - k > \text{ord}_p d! - \lfloor \log_p d \rfloor$, will satisfy inequality $k \geq \lambda(f)$. Yet since $\text{ord}_p d! = \frac{1}{p-1}(d - \text{wt}_p d)$, where $\text{wt}_p d$ is a sum of all digits in p -base expansion of d , then, choosing any $k \in \mathbb{N}$, which satisfy inequality

$$2 \frac{p^k - 1}{p - 1} - k > \frac{d}{p - 1}, \quad (1)$$

we obtain that $k \geq \lambda(f)$. Elementary considerations, however, show that $k = \lfloor \log_p d \rfloor + 1$ satisfies inequality (1), thus proving the proposition for $p \neq 2$.

In case $p = 2$ a polynomial $f(x) \in \mathbb{Q}_2[x]$ of degree d is integer-valued, compatible and ergodic iff it is of a form

$$f(x) = 1 + x + \sum_{i=0}^d b_i 2^{\lfloor \log_2(i+1) \rfloor + 1} \binom{x}{i}, \quad (2)$$

where $b_i \in \mathbb{Z}_2$, $i = 0, 1, 2, \dots, d$ (see theorem 2.3). Since coefficients of the polynomial $f(x)$ in its representation in a basis $\binom{x}{i}$, $i = 0, 1, 2, \dots$, are uniquely defined by the values of $f(z)$ at the points $z = 0, 1, \dots, d$, then to verify conditions (2) for the polynomial $f(x)$ it is sufficient to calculate its values at the points $z = 0, 1, \dots, 2^r - 1$, where $r \in \mathbb{N}$ is an arbitrarily fixed number satisfying the inequality $d \leq 2^r - 1$. So one can take, for instance, $r = \lfloor \log_2(d + 1) \rfloor + 1$, or $r = \lfloor \log_2 d \rfloor + 3$. This finishes the proof of 4.7. $\square$

Note. Proposition 4.4 shows that for $p \neq 2$ a function $f \in \mathcal{A}$ satisfies assumptions of the proposition 3.9; hence, since $N_1(f) \leq N_2(f)$, a function f preserves measure iff it is bijective modulo $p^{\lambda(f)+2}$. By the argument similar to those of the proof of proposition 4.7, one could prove the following

4.8 Proposition. *A polynomial $f(x) \in \mathbb{Q}_p[x]$ is integer-valued, compatible and measure-preserving iff the mapping*

$$z \mapsto f(z) \bmod p^{K_f},$$

with $K_f = \lfloor \log_p(\deg f) \rfloor + 3$ and z ranging over $0, 1, \dots, p^{K_f} - 1$, induces a compatible and bijective function on the ring $\mathbb{Z}/p^{K_f}$. $\square$

Again, estimates of $\zeta(f)$ and $\eta(f)$ we mentioned at the beginning of the section, might be sharpened for various important proper subclasses of $\mathcal{A}$ in comparison

with given by the theorem 4.1 and propositions 4.7 and 4.8. A case of analytic on $\mathbb{Z}_p$ functions (i.e., functions which can be represented by convergent everywhere on $\mathbb{Z}_p$ power series) seems to be of importance.

It is well known (see e.g. [3, Ch. 14. Section 4]) that power series $\sum_{i=0}^{\infty} c_i x^i$ ($c_i \in \mathbb{Q}_p$, $i = 0, 1, 2, \dots$) converges everywhere on $\mathbb{Z}_p$ iff $\lim_{i \rightarrow \infty}^p c_i = 0$; under the latter condition the series defines a continues function on $\mathbb{Z}_p$. Of course, in general this function may not be integer-valued, not speaking about compatibility. Consider, however, a particular case, when all coefficients c_i are p -adic integers. Namely, in the ring $\mathbb{Z}_p[[x]]$ of all formal power series in variable x over a ring $\mathbb{Z}_p$ consider a set $\mathcal{C}(x)$ of all series

$$s(x) = \sum_{i=0}^{\infty} c_i x^i \quad (c_i \in \mathbb{Z}_p, i = 0, 1, 2, \dots),$$

which converge everywhere on $\mathbb{Z}_p$. In other words, $s(x) \in \mathcal{C}(x)$ iff $\lim_{i \rightarrow \infty}^p c_i = 0$.

Under these assumptions the series $s(x) \in \mathcal{C}(x)$ defines on $\mathbb{Z}_p$ an integer-valued function $s : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$. It turns out that this function s is uniformly differentiable and has integer-valued derivative everywhere on $\mathbb{Z}_p$.

Consider a formal derivative $s'(x) \in \mathbb{Z}_p[[x]]$ of the series $s(x)$:

$$s'(x) = \sum_{i=1}^{\infty} i c_i x^{i-1}.$$

Since $0 \leq \|i c_i\|_p = \|i\|_p \|c_i\|_p \leq \|c_i\|_p$, and $\lim_{i \rightarrow \infty}^p c_i = 0$, then $\lim_{i \rightarrow \infty}^p i c_i = 0$, and hence $s'(x) \in \mathcal{C}(x)$. We assert that the function $s' : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is a derivative of a function $s : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ with respect to p -adic distance.

Indeed, it is known that in the ring $\mathbb{Z}_p[[x, y]]$ of all formal power series in variables x, y over $\mathbb{Z}_p$ the following equality holds:

$$s(x + y) = \sum_{i=0}^{\infty} \frac{s^{(i)}(x)}{i!} y^i,$$

where $s^{(i)}(x) \in \mathbb{Z}_p[[x]]$ ($i = 1, 2, \dots$) is i th formal derivative of the series $s(x)$, and $s^{(0)}(x) = s(x)$. By the assertion proven above, $s^{(i)}(x) \in \mathcal{C}(x)$ for all $i = 0, 1, 2, \dots$. Thus,

$$\frac{s^{(i)}(u)}{i!} = \sum_{j=i}^{\infty} c_j \binom{j}{i} u^{j-i} \in \mathbb{Z}_p$$

for each $u \in \mathbb{Z}_p$. But

$$\left\| \frac{s^{(i)}(u)}{i!} \right\|_p = \left\| \sum_{j=i}^{\infty} c_j \binom{j}{i} u^{j-i} \right\|_p \leq \max\{\|c_j\|_p : j = i, i+1, \dots\},$$

and consequently,

$$\lim_{i \rightarrow \infty}^p \frac{s^{(i)}(u)}{i!} = 0,$$

since $\lim_{i \rightarrow \infty}^p c_i = 0$. Thus, for each $u \in \mathbb{Z}_p$ we have that

$$s(u + y) = \sum_{i=0}^{\infty} \frac{s^{(i)}(u)}{i!} y^i \in \mathcal{C}(y). \quad (\spadesuit)$$

Finally, if $s(x) \in \mathcal{C}(x)$, then Taylor series $(\spadesuit)$ at the point $u \in \mathbb{Z}_p$ converges to s everywhere on $\mathbb{Z}_p$. In particular, for $h \in \mathbb{Z}_p$ we obtain

$$s(u + h) = s(u) + s'(u)h + \alpha(u, h),$$

with $\lim_{h \rightarrow 0}^p \frac{\alpha(u, h)}{h} = \lim_{h \rightarrow 0}^p h \sum_{i=2}^{\infty} \frac{s^{(i)}(u)}{i!} h^{i-2} = 0$, since $\sum_{i=2}^{\infty} \frac{s^{(i)}(u)}{i!} h^{i-2} \in \mathbb{Z}_p$ in view of the equality $\lim_{i \rightarrow \infty}^p \frac{s^{(i)}(u)}{i!} = 0$, which just has been proven above. So, $s'(u)$ is a derivative of the function s at the point u . Thus, the set $\mathcal{C}(x)$ is closed with respect to differentiations, and all functions, defined by series of $\mathcal{C}(x)$, are infinitely many times differentiable.

Further, let

$$s(x) = \sum_{i=0}^{\infty} s_i \binom{x}{i}$$

be an interpolation series for the function $s(x) \in \mathcal{C}(x)$. We assert that $\frac{s_i}{i!}$ is p -adic integer for all $i = 0, 1, 2, \dots$. Actually,

$$s(x) = \sum_{k=0}^{\infty} c_k x^k = \sum_{k=0}^{\infty} c_k \sum_{i=0}^k S_2(k, i) i! \binom{x}{i} = \sum_{i=0}^{\infty} i! \binom{x}{i} \sum_{k=i}^{\infty} S_2(k, i) c_k,$$

where $S_2(k, i)$ is Stirling number. Since $\lim_{i \rightarrow \infty}^p c_i = 0$, then $\lim_{k \rightarrow \infty}^p S_2(k, i) c_k = 0$, because all Stirling numbers $S_2(k, i)$ are integer rationals, i.e., $\|S_2(k, i)\|_p \leq 1$. Consequently, the series $\sum_{k=i}^{\infty} S_2(k, i) c_k$ converges to some $A_i \in \mathbb{Z}_p$ for all $i = 0, 1, 2, \dots$. This proves our assertion, since

$$s_i = i! A_i \quad (i = 0, 1, 2, \dots). \quad (\star)$$

Put

$$\mathcal{B}(x) = \left\{ f(x) = \sum_{i=0}^{\infty} a_i \binom{x}{i} : \frac{a_i}{i!} \in \mathbb{Z}_p, \quad i = 0, 1, 2, \dots \right\}.$$

In other words, $\mathcal{B}(x)$ is a ring of all formal descending factorial power series over $\mathbb{Z}_p$. Each series $f(x) \in \mathcal{B}(x)$ correctly defines on $\mathbb{Z}_p$ an integer-valued and uniformly continuous function $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ (see the beginning of the section 2). This function f is compatible in view of 2.1, since we have shown during the proof of the lemma 4.2 that $\text{ord}_p(i!) - \lfloor \log_p i \rfloor$ is nonnegative and nondecreasing function on $\mathbb{N}_0$. Denote via $\mathcal{B}$ (respectively, via $\mathcal{C}$) a class of all functions defined by all series of $\mathcal{B}(x)$ (respectively, of $\mathcal{C}(x)$). Obviously, $\mathcal{B}(x)$, $\mathcal{B}$, $\mathcal{C}(x)$, $\mathcal{C}$ are rings.

Further, any two distinct series of $\mathcal{B}(x)$ (respectively, of $\mathcal{C}(x)$) define two distinct functions on $\mathbb{Z}_p$: for the series of $\mathcal{B}(x)$ see the beginning of the section 2. As for

the series of $\mathcal{C}(x)$, note, that the above mentioned interpolation series for $s(x) \in \mathcal{C}(x)$ defines a function, which is identically 0 on $\mathbb{Z}_p$ iff all its coefficients s_i are 0 (hence, $A_i = 0$, $i = 0, 1, 2, \dots$), see (★). Yet $A_i = \sum_{k=i}^{\infty} S_2(k, i)c_k$, hence $c_i = \sum_{k=i}^{\infty} S_1(k, i)A_k = 0$, where $S_1(k, i), S_2(k, i)$ are Stirling numbers of respective kind, and the assertion follows. Thus, the rings $\mathcal{B}(x)$ and $\mathcal{B}$ (respectively, $\mathcal{C}(x)$ and $\mathcal{C}$) are isomorphic; so further we do not differ series from the function it defines.

Note also that the inclusion $\mathcal{B} \supset \mathcal{C}$ (see (★)) is strict. Obviously, $f(x) = \sum_{i=0}^{\infty} (x)_i \in \mathcal{B}$, since $f(x) = \sum_{i=0}^{\infty} i! \binom{x}{i}$. Yet $f(x) \notin \mathcal{C}$. Moreover, this function is not even analytic on $\mathbb{Z}_p$: according to [3, Ch.4, Theorem 4] a function represented by the interpolation series ($\diamond$) of the section 2 is analytic on $\mathbb{Z}_p$ iff $\lim_{i \rightarrow \infty} \frac{a_i}{i!} = 0$.

So, a function of $\mathcal{B}$ (in contrast to one of $\mathcal{C}$), generally speaking, can not be represented by Taylor series which is convergent everywhere on $\mathbb{Z}_p$. Nevertheless, all functions of $\mathcal{B}$ are differentiable at all points of $\mathbb{Z}_p$, and $\mathcal{B}$ is closed with respect to differentiations: if $f \in \mathcal{B}$, then $f' \in \mathcal{B}$.

To prove the latter assertion, recall that a uniformly continuous on $\mathbb{Z}_p$ function f , which is represented by the interpolation series ($\diamond$), is differentiable everywhere on $\mathbb{Z}_p$ iff

$$\lim_{i \rightarrow \infty} \frac{a_{i+n}}{i} = 0 \quad (\diamond)$$

for all $n \in \mathbb{N}_0$ (see [3, Ch. 13, Theorem 2]). The latter condition obviously holds for $f \in \mathcal{B}$, since $\text{ord}_p a_i \geq \text{ord}_p (i!) = \frac{1}{p-1}(i - \text{wt}_p i)$, and $\lfloor \log_p i \rfloor \geq \text{ord}_p i$ for all $i = 0, 1, 2, \dots$. Thus, a derivative f' of the function f is defined everywhere on $\mathbb{Z}_p$, and

$$f'(x) = \sum_{i=1}^{\infty} (-1)^{i+1} \frac{\Delta^i f(x)}{i},$$

in case this series is convergent. Yet $\frac{\Delta^i f(x)}{i} = \frac{1}{i} \sum_{j=i}^{\infty} a_j \binom{x}{j-i}$, consequently,

$$\sum_{i=1}^{\infty} (-1)^{i+1} \frac{\Delta^i f(x)}{i} = \sum_{k=0}^{\infty} \binom{x}{k} \sum_{i=1}^{\infty} (-1)^{i+1} \frac{a_{k+i}}{i}.$$

But the series $\sum_{i=1}^{\infty} (-1)^{i+1} \frac{a_{k+i}}{i}$, in view of ($\diamond$), for each $k \in \mathbb{N}_0$ converges to a certain $S_k \in \mathbb{Q}_p$, and $\text{ord}_p \frac{a_{k+i}}{i} = \text{ord}_p a_{k+i} - \text{ord}_p i \geq \text{ord}_p ((k+i)!) - \lfloor \log_p i \rfloor = \frac{1}{p-1}(i + k - \text{wt}_p(i+k)) - \lfloor \log_p i \rfloor = \frac{1}{p-1}(i - \text{wt}_p i) - \lfloor \log_p i \rfloor + \frac{1}{p-1}(k - \text{wt}_p k) + \frac{1}{p-1}(\text{wt}_p k - \text{wt}_p(i+k) + \text{wt}_p i) \geq \frac{1}{p-1}(k - \text{wt}_p k) = \text{ord}_p(k!)$. (The latter inequality holds since $\frac{1}{p-1}(i - \text{wt}_p i) \geq \lfloor \log_p i \rfloor$ and $\frac{1}{p-1}(\text{wt}_p k - \text{wt}_p(i+k) + \text{wt}_p i) = \text{ord}_p \binom{i+k}{i} \geq 0$).

Thus, $\frac{S_k}{k!} \in \mathbb{Z}_p$ for all $k \in \mathbb{N}_0$; hence $f' \in \mathcal{B}$.

With the use of these results now we are able to prove the following

4.9 Theorem. *A function $f \in \mathcal{B}$ preserves measure iff it is bijective modulo p^2 . The function f is ergodic iff it is transitive modulo p^2 (for $p \neq 2, 3$), or modulo p^3 (for $p \in \{2, 3\}$).*

Proof. The definition of $\mathcal{B}$ immediately implies that $\rho(f) = 0$ for each $f \in \mathcal{B}$, hence, $\lambda(f) = 1$. Thus, for $p \neq 2$ the second assertion of the theorem follows from 4.1.

To prove the first assertion, in view of 3.9 it is sufficient to demonstrate that f is uniformly differentiable modulo p , and $N_1(f) \leq 1$; that is

$$f(z + p^k r) \equiv f(z) + p^k r f'(z) \pmod{p^{k+1}} \quad (1)$$

for all $z, r \in \mathbb{Z}_p$ and $k = 1, 2, \dots$. Since $f, f' \in \mathcal{B}$, these both functions are compatible, so it is sufficient to prove (1) for $z, r \in \mathbb{N}_0$. Since for $r = 0$ the congruence (1) is trivial, we may additionally assume that $p^k r = n \in \mathbb{N}$.

Further, since $\frac{f(z+n)-f(z)}{n} = \sum_{i=1}^{\infty} \binom{n-1}{i-1} \frac{\Delta^i f(x)}{i}$, $f'(z) = \sum_{i=1}^{\infty} (-1)^{i+1} \frac{\Delta^i f(z)}{i}$, then to prove (1) it is sufficient to prove that

$$\sum_{i=1}^{\infty} \left(\binom{n-1}{i-1} - (-1)^{i+1} \right) \frac{\Delta^i f(z)}{i} \equiv 0 \pmod{p}. \quad (2)$$

Yet $\frac{\Delta^i f(x)}{i} = \frac{1}{i} \sum_{j=i}^{\infty} a_j \binom{x}{j-i}$, thus, in view of 4.2, for $p \neq 2$ there holds a congruence $\frac{\Delta^i f(x)}{i} \equiv 0 \pmod{p}$ for all $i \geq 2p$. So within this case (2) is equivalent to the congruence

$$\sum_{i=1}^{2p-1} \left(\binom{n-1}{i-1} - (-1)^{i+1} \right) \frac{\Delta^i f(z)}{i} \equiv 0 \pmod{p}. \quad (3)$$

Since f is compatible, then $\frac{\Delta^i f(x)}{i} \not\equiv 0 \pmod{p}$ only for, might be, $i = sp^m$, ($m \in \mathbb{N}_0$, $s \in \{1, 2, \dots, p-1\}$) — see [11, lemma 3.4]. Now, since $n = p^k r$, (3) immediately follows from the already mentioned Lucas theorem, thus proving the first assertion of 4.9 for $p \neq 2$.

Now, if $p = 2$, then (2) is equivalent to

$$\sum_{i=0}^{\infty} \left(\binom{2^k r - 1}{2^i - 1} + 1 \right) \frac{\Delta^{2^i} f(z)}{2^i} \equiv 0 \pmod{2}. \quad (4)$$

Yet since $\frac{a_j}{j!} \in \mathbb{Z}_2$ for all $j = 0, 1, 2, \dots$, then $\text{ord}_2 a_{2^i+m} \geq \text{ord}_2 (2^i)! = 2^i - 1$ for all $m = 0, 1, 2, \dots$; consequently, $\frac{\Delta^{2^i} f(z)}{2^i} \not\equiv 0 \pmod{2}$ only for, might be, $i = 0$, thus proving (4).

Finally, the rest part of the assertion of theorem 4.9 for $p = 2$ follows from 2.3: as $\text{ord}_2 i! \leq \text{ord}_2 a_i$ for all $i = 0, 1, 2, \dots$, and $\text{ord}_2 i! = i - \text{wt}_2(i)$, then by an elementary argument it is not difficult to show that $\lfloor \log_2(i+1) \rfloor + 1 \leq i - \text{wt}_2(i) \leq \text{ord}_2 a_i$ for $i \geq 4$; and $\text{ord}_2 a_i \geq 3$. This implies that necessary and sufficient conditions of ergodicity of a function expressed as interpolation series ($\diamond$) of section 2 hold for all coefficients a_i with $i \geq 4$. These conditions for the rest of the coefficients are equivalent to the transitivity of f modulo 8, since $a_i \equiv 0 \pmod{8}$ for $i \geq 4$. $\square$

Note. Theorem 4.9 demonstrates that sufficient and necessary conditions of transitivity modulo p^n for the polynomials with integer rational coefficients established by M. V. Larin in [15] remain valid for a wider class (namely, $\mathcal{B}$) of functions. It turns out, however, that all these functions modulo each p^n could be expressed as polynomials with rational integer coefficients.

Namely, from the definition of a class $\mathcal{B}$ it easily follows that each function $f \in \mathcal{B}$ is uniformly approximated by polynomials over $\mathbb{Z}_p$: for each $n \in \mathbb{N}$ there

exists a polynomial $f_n(x) \in \mathbb{Z}_p[x]$, such that $f(z) \equiv f_n(z) \pmod{p^n}$ for all $z \in \mathbb{Z}_p$. Actually, the series $\sum_{j=0}^{\infty} r_j \binom{x}{j}$ defines a function, which is identically 0 modulo p^n iff all $r_j \equiv 0 \pmod{p^n}$ (see [11, proposition 4.2]). So we may put $f_n(x) = \sum_{i=0}^{\omega(n)} a_i \binom{x}{i}$, where $\omega(n) = \max\{j \in \mathbb{N}_0 : \frac{1}{p-1}(j - \text{wt}_p j) < n\}$.

It turns out that the inverse assertion is also true: if a function $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is uniformly approximated by polynomials over $\mathbb{Z}_p$ in the above mentioned sense, then it lies in $\mathcal{B}$. To prove this assertion, assume that $f(z) \equiv f_i(z) \pmod{p^i}$ for all $z \in \mathbb{Z}_p$, where $f_i(x) \in \mathbb{Z}_p[x]$, $i = 1, 2, \dots$. Each polynomial $f_i(x)$ of degree d_i admits one and the only representation as interpolation series ($\diamond$) of section 2: $f_i(x) = \sum_{j=0}^{d_i} a_{ij} \binom{x}{j}$, where $a_{ij} \in \mathbb{Z}_p$ and $\text{ord}_p a_{ij} \geq \text{ord}_p(j!)$ in view of ($\star$), since, obviously, $f_i \in \mathcal{C} \subset \mathcal{B}$. For a given function f each polynomial $f_i(x)$ is unique up to the summand which induces an identically 0 modulo p^i function. So we may assume that $d_i = \omega(i)$ (see above); then coefficients of the polynomial $f_i(x)$ are defined uniquely up to the summands with p -adic norms not exceeding p^{-i} . This implies that $a_{i+1,j} \equiv a_{ij} \pmod{p^i}$ (we assume $a_{ij} = 0$ for $j > \omega(i)$). Hence, $\lim_{i \rightarrow \infty}^p a_{ij} = a_j \in \mathbb{Z}_p$, and $\frac{a_j}{j!} \in \mathbb{Z}_p$. Consequently, the series $\sum_{i=0}^{\infty} a_i \binom{x}{i}$ defines a uniformly continuous on $\mathbb{Z}_p$ function $\tilde{f} \in \mathcal{B}$, which must be equal to f , since $f(z) \equiv f_i(z) \equiv \tilde{f}(z) \pmod{p^i}$ for all $z \in \mathbb{Z}_p$ and all $i = 1, 2, \dots$.

Now we define a non-Archimedean pseudo-valuation on $\mathcal{B}$ as $\max\{\|f(z)\|_p : z \in \mathbb{Z}_p\}$ for $f \in \mathcal{B}$. The just proven results imply that with respect to the distance D_p , induced by this pseudo-valuation, the ring $\mathcal{B}$ is a complete metric space; actually, $\mathcal{B}$ is a completion with respect to D_p of the space $\mathcal{P} \subset \mathcal{C}$ of all functions induced on $\mathbb{Z}_p$ by polynomials over $\mathbb{Z}$ (in particular, the space $\mathcal{B}$ is separable).

This implies, in turn, that $\mathcal{B}$ (contrasting to $\mathcal{C}$) is closed with respect to composition of functions: if $f, g \in \mathcal{B}$ then $f(g) \in \mathcal{B}$. In fact, let g be uniformly approximated by the sequence $\{g_n(x) \in \mathbb{Z}_p[x] : n = 1, 2, \dots\}$, that is, $g_n(z) \equiv g(z) \pmod{p^n}$ for all $z \in \mathbb{Z}_p$. The compatibility of the function f implies then that $D_p(f(g), f(g_n)) \leq p^{-n}$, i.e., for $n \rightarrow \infty$ the sequence $f(g_n)$ tends to $f(g)$ with respect to distance D_p . But $f(g_n) \in \mathcal{B}$ for each $n = 1, 2, \dots$: if f is uniformly approximated by the sequence $\{f_m(x) \in \mathbb{Z}_p[x] : m = 1, 2, \dots\}$, then $f_m(g_n(z)) \equiv f(g_n(z)) \pmod{p^m}$ for all $z \in \mathbb{Z}_p$. Hence, the sequence $\{f_m(g_n(x)) \in \mathbb{Z}_p[x] : m = 1, 2, \dots\}$ tends to the function $f(g_n)$ with respect to the distance D_p , and $f_m(g_n) \in \mathcal{B}$, since it is a polynomial over $\mathbb{Z}_p$. Consequently, $f(g) \in \mathcal{B}$ in view of completeness of $\mathcal{B}$.

Thus, we have proven the following

4.10 Proposition. *The ring $\mathcal{B}$ is a separable and complete with respect to the distance D_p metric space of functions, which are differentiable everywhere on $\mathbb{Z}_p$. $\mathcal{B}$ is closed with respect to compositions of functions and with respect to differentiations. A countable set $\mathcal{P}$ of all polynomials over $\mathbb{Z}$ is a dense subset of $\mathcal{B}$. $\square$*

To make use of criterion 4.9 for the applications to pseudorandom number generation it is important to have a huge stock of examples of functions of $\mathcal{B}$ which are to be implemented as computer programs. As we have mentioned above, all polynomials over $\mathbb{Z}_p$ are in $\mathcal{B}$.

Rational over $\mathbb{Z}_p$ functions, that is, functions of the form $f(x) = \frac{u(x)}{v(x)}$, where $u(x), v(x) \in \mathbb{Z}_p[x]$, are also in $\mathcal{B}$, providing the denominator vanishes modulo p nowhere on $\mathbb{Z}_p$ (in view of compatibility it is sufficient to verify the latter condition only for the points of $\{0, 1, \dots, p-1\}$). Indeed, for each $z \in \mathbb{Z}_p$ the element

$v(z)$ is not 0 modulo p , and hence has a multiplicative inverse in the ring $\mathbb{Z}/p^n$. Thus $\frac{u(z)}{v(z)} \equiv u(z)v(z)^{\phi(p^n)-1} \pmod{p^n}$, where ϕ is Euler totient function. Hence, the function f could be uniformly approximated by polynomials $u(x)v(x)^{\phi(p^n)-1} \in \mathbb{Z}_p[x]$, $n = 1, 2, \dots$; hence, it is in $\mathcal{B}$ in force of 4.10.

Another type of functions of $\mathcal{B}$ are exponential ones. For instance, consider a function a^x with $a \equiv 1 \pmod{p}$ (hence, $a = 1 + pr$ for suitable $r \in \mathbb{Z}_p$). Then $a^x = \sum_{i=0}^{\infty} p^i r^i \binom{x}{i}$, and it is well known (see e.g. [3, Ch. 14, Section 5]), that for $p \neq 2$ this function is analytic on $\mathbb{Z}_p$ (hence, lies in $\mathcal{C}$). If $p = 2$ and r is odd, then a^x is not analytic on $\mathbb{Z}_2$, thus not in $\mathcal{C}$. Nevertheless, within the latter case a^x is in $\mathcal{B}$, since $\text{ord}_2(i!) = i - wt_2 i$ and hence $(1 + 2r)^x = \sum_{i=0}^{\infty} 2^i r^i \binom{x}{i} \in \mathcal{B}$. It is not difficult to see that the function $(1 + 4r)^x$ is in $\mathcal{C}$. So, summarizing all these considerations, if $a \in \mathbb{Z}_p$, $a \equiv 1 \pmod{p}$, then the function a^x is in $\mathcal{B}$.

Exponential functions of the considered type are particular cases of functions of more general form u^v , where $u(z) \equiv 1 \pmod{p}$ for all $z \in \mathbb{Z}_p$.

4.11 Lemma. *Let $u, v: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be compatible functions and let $u(z) \equiv 1 \pmod{p}$ for all $z \in \mathbb{Z}_p$ (so it is sufficient to verify the latter condition only for $z = 0, 1, \dots, p-1$). Then the function $f(z) = u(z)^{v(z)}$ is correctly defined for all $z \in \mathbb{Z}_p$, integer-valued and compatible. Moreover, if $w, v \in \mathcal{B}$, $u(z) = 1 + pw(z)$, then $f \in \mathcal{B}$.*

Proof. The above considerations of functions of type a^x with $a \equiv 1 \pmod{p}$ immediately imply that the function f is correctly defined on $\mathbb{Z}_p$ and that it is integer-valued. To prove the compatibility of f , note, that for arbitrary $b, c, d \in \mathbb{Z}_p$ and $n = 1, 2, \dots$ one has $(a + p^n b)^{c+p^n d} = (a + p^n b)^c ((a + p^n b)^{p^n})^d$, since elementary properties of powers are of the same form both in real and p -adic cases, see [3, Ch. 14, Section 5]. As both u and v are compatible functions, then for arbitrary $z, r \in \mathbb{Z}_p$ there exist $s, t \in \mathbb{Z}_p$, such that $(u(z + p^n r))^{v(z + p^n r)} = (u(z) + p^n t)^{v(z) + p^n s}$; hence $(u(z + p^n r))^{v(z + p^n r)} = (u(z) + p^n t)^{v(z)} ((u(z) + p^n t)^{p^n})^s \equiv (u(z) + p^n t)^{v(z)} \pmod{p^n}$, in view of the congruence $(u(z) + p^n t)^{p^n} \equiv 1 \pmod{p^n}$. The latter congruence is to be proven.

As $u(z) \equiv 1 \pmod{p}$, then for a suitable $k \in \mathbb{Z}_p$ we have $u(z) + p^n t = 1 + pk$. Yet $(1 + pk)^{p^n} = \sum_{i=0}^{p^n} k^i p^i \binom{p^n}{i} = \sum_{i=0}^{p^n} k^i \frac{p^i}{i!} (p^n)_i \equiv 1 \pmod{p^n}$, since $\frac{p^i}{i!} \in \mathbb{Z}_p$. Finally, denoting by $\overline{v(z)} = v(z) \bmod p^n$ the least nonnegative residue of $v(z)$ modulo p^n , for a suitable $h \in \mathbb{Z}_p$ we obtain $f(z + p^n r) \equiv (u(z) + p^n t)^{v(z)} = (u(z) + p^n t)^{\overline{v(z)}} (u(z) + p^n t)^{p^n h} \equiv (u(z) + p^n t)^{\overline{v(z)}} = \sum_{i=0}^{\overline{v(z)}} u(z)^{\overline{v(z)}-i} p^{ni} t^i \binom{\overline{v(z)}}{i} = (u(z))^{\overline{v(z)}} \equiv (u(z))^{\overline{v(z)}} (u(z))^{p^n h} = (u(z))^{v(z)}$, where $\equiv$ stands for congruence modulo p^n . Thus, f is compatible.

To prove the rest of the lemma, note, that for each $z \in \mathbb{Z}_p$ and each $n = 1, 2, \dots$ the congruence $(u(z))^{v(z)} \equiv \sum_{i=0}^n (u(z) - 1)^i \binom{v(z)}{i} \pmod{p^n}$ holds, since $\|u(z) - 1\|_p \leq \frac{1}{p}$. This implies that

- (1) all functions $f_n = \sum_{i=0}^n \frac{p^i}{i!} (v)_i w^i$ are in $\mathcal{B}$, since all $\frac{p^i}{i!}$ are p -adic integers (see above);
- (2) the sequence $\{f_n : n = 1, 2, \dots\}$ tends to f with respect to the distance D_p .

Now (1)–(2) imply that $f \in \mathcal{B}$ in force of 4.10. $\square$

With the use of these results one may construct explicit forms of various ergodic functions to be performed by a computer. For instance, the following is true.

4.12 Proposition. For $g \in \mathcal{B}$ the function $f(x) = 1 + x + p^2 g(x)$ is ergodic.

Proof. For $p \notin \{2, 3\}$ the assertion trivially follows from 4.9. For $p \in \{2, 3\}$ in view of 4.9 it is sufficient to show that f is transitive modulo p^3 . In turn, to demonstrate the latter it is sufficient to prove only that $f^{kp^2}(0) \not\equiv 0 \pmod{p^3}$ for $k = 1, 2, \dots, p-1$, since in f is transitive modulo p^2 and hence, in view of its compatibility, induces on $\mathbb{Z}/p^3$ a permutation with each cycle length being a multiple of p^2 . Yet since for all $i = 0, 1, 2, \dots$ the compatibility of g implies that $f^i(0) \equiv i + p^2 \sum_{j=0}^{i-1} g(j) \pmod{p^3}$, then $f^{kp^2}(0) \equiv kp^2 + p^2 \sum_{j=0}^{kp^2-1} g(j) \equiv kp^2 + p^2 \sum_{z=0}^{p-1} g(z)pk \equiv kp^2 \pmod{p^3}$, since (again in view of the compatibility of g) a congruence $s \equiv r \pmod{p}$ implies the congruence $p^2 g(r) \equiv p^2 g(s) \pmod{p^3}$. $\square$

5. APPLICATIONS: A DISCUSSION.

The results obtained in previous sections might have applications to design pseudorandom number generators which have relatively simple program implementation, generate purely periodic sequences of numbers of $\{0, 1, \dots, m-1\}$ and provide certain guarantee for the statistical quality of these sequences, their uniform distribution at the first turn. Speaking about relatively simple program implementation, we mean that the considered generators have certain parameters which are critical to the performance, and which one may vary to achieve the desired performance without affecting the quality.

In case $m = p^k$ is a power of a prime p , these sequences might be generated as the first order recurrence sequences satisfying the relation $x_{n+1} \equiv f(x_n) \pmod{m}$, where $f: \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ is any compatible and ergodic function of the considered in previous sections. In this case for each $k = 1, 2, \dots$ we obtain a purely periodic sequence with period length p^k , with each element of $\{0, 1, \dots, p^k-1\}$ occurring at the period exactly once (in particular, the generated sequence is uniformly distributed).

An important indicator of statistical quality of the sequence is the distribution of $(r+1)$ -tuples $\{(x_n, x_{n+1}, \dots, x_{n+r}) : n = 0, 1, 2, \dots\}$. Ideally, the sequence $\{\mathbf{u}_n = (\frac{x_n}{p^k}, \frac{x_{n+1}}{p^k}, \dots, \frac{x_{n+r}}{p^k}) : n = 0, 1, 2, \dots\}$ of points of $(r+1)$ -dimensional Euclidean space should be uniformly distributed in the unit hypercube for all r . By no means this can be achieved for periodic sequences. For such sequences there are some popular tests of quality, based on certain characteristics of families of hyperplanes, which are parallel one to another, and which union contain all the points corresponding to the sequences of $(r+1)$ -tuples (see e.g. [2, section 3.3.4]).

Note, that if for some $c, c_0, \dots, c_r \in \mathbb{Z}$ the congruences

$$c + \sum_{i=0}^r c_i x_{n+i} \equiv 0 \pmod{p^k}, \quad (n = 0, 1, 2, \dots) \quad (\blacktriangle)$$

hold, then all the points $\mathbf{u}_n$ fall into the hyperplanes $h + \sum_{i=0}^r c_i X_i = 0$, which are parallel one to another. For linear congruential generators such families of parallel hyperplanes exist even for $r = 2$, not depending on k (see the introduction).

Note, that if $(\blacktriangle)$ holds for some k , then for all $j = 1, 2, \dots$ for the members of the sequence $\{x_n\}$ hold relations $p^j c + \sum_{i=0}^r p^j c_i x_{n+i} \equiv 0 \pmod{p^{k+j}}$. The relations of the latter kind will be temporarily and loosely defined as trivial. Trivial relations always exist: for instance, choosing certain $K \in \mathbb{N}$ in view of the ergodicity of f we obtain for all $k \geq K$ the trivial relations $p^{k-K} x_{n+p^K} \equiv p^{k-K} x_n \pmod{p^k}$.

Speaking informally, the triviality of relations just means that their coefficients tend to 0 whereas k tends to infinity, i.e. trivial relations are those which degenerate to $0 = 0$ in $\mathbb{Z}_p$.

For an important wide class of nonlinear congruential generators we prove that if the dimension of hyperplanes, which are parallel one to another, and which union contains all points $\mathbf{u}_n$, ($n = 0, 1, 2, \dots$), does not tend to infinity together with k , then this family of hyperplanes is defined by trivial relations.

Now we give exact statements.

5.1 Proposition. *Let $f \in \mathbb{Q}_p[x]$ be an integer-valued, compatible and ergodic polynomial of degree d over a field $\mathbb{Q}_p$ of p -adic numbers (all these polynomials for $p = 2$ are completely characterized by theorem 2.3; for odd p see 2.4, 4.7 and a note preceding 4.7). Let, further, r be a positive integer rational such that for each $k \in \mathbb{N}$ there exist $c, c_0, \dots, c_r \in \mathbb{Z}_p$, which satisfy $(\blacktriangle)$ and not all of which are 0 modulo p . Then $d = 1$.*

We will need the following

5.2 Lemma. *Under the assumptions of proposition 5.1 let $c, c_0, \dots, c_r \in \mathbb{Z}_p$ be not depending on k , that is, let there exist $c, c_0, \dots, c_r \in \mathbb{Z}_p$ satisfying $(\blacktriangle)$ for all $k \in \mathbb{N}$ simultaneously. Then $d = 1$.*

Proof of the lemma 5.2. As f is ergodic, then $d \neq 0$. Assume that $d > 1$. Consider $w(x) = c + \sum_{i=0}^r c_i f^i(x)$. As $w(x)$ is a composition of integer-valued and compatible polynomials over $\mathbb{Q}_p$, then $w(x) \in \mathbb{Q}_p[x]$ is integer-valued and compatible. Yet each $f^i(x)$ has degree d^i ; hence, since $d > 1$, then $w(x)$, being a sum of polynomials of pairwise distinct degrees, must be a polynomial of nonzero degree.

Yet, since $x_{n+i} \equiv f^i(f^n(x_0)) \pmod{p^k}$, the assumptions of the lemma imply that $w(x_n) \equiv 0 \pmod{p^k}$ for all $n = 0, 1, 2, \dots$. In other words, $w(z) \equiv 0 \pmod{p^k}$ for all $z \in \mathbb{Z}_p$, since x_n takes all values in $\{0, 1, \dots, p^k - 1\}$ in view of the ergodicity of f , and $w(x)$ is compatible. The assumptions of the lemma now imply that $w(z) \equiv 0 \pmod{p^k}$ for all $z \in \mathbb{Z}_p$ and all $k = 1, 2, \dots$. Consequently, $w(z) = 0$ for all $z \in \mathbb{Z}_p$ and hence polynomial $w(x)$ must be 0 in the ring $\mathbb{Q}_p[x]$. A contradiction proving the lemma. $\square$

Proof of the proposition 5.1. By the assumption, for each $k \in \mathbb{N}$ the set $\mathcal{L}_k$ of all $\mathbf{c} = (c, c_0, \dots, c_r) \in \mathbb{Z}_p^{r+2}$, $\|\mathbf{c}\|_p = 1$ with $c, c_0, \dots, c_r$ satisfying $(\blacktriangle)$, is not empty. Obviously, $\mathcal{L}_1 \supset \mathcal{L}_2 \supset \dots$, since f is compatible.

Further, we assert that each set $\mathcal{L}_k$ is closed in the topology of metric space $\mathbb{Z}_p^{r+2}$. Actually, if $\mathbf{c} \in \mathcal{L}_k$, $\mathbf{c}' \in \mathbb{Z}_p^{r+2}$, $\|\mathbf{c} - \mathbf{c}'\| \leq p^{-s}$, $s \geq k$, then $\mathbf{c}' = \mathbf{c} + p^s \mathbf{z}$ for a suitable $\mathbf{z} \in \mathbb{Z}_p^{r+2}$. Hence, $\|\mathbf{c}'\|_p = 1$ and $\mathbf{c}'$ satisfies $(\blacktriangle)$; consequently, $\mathbf{c}' \in \mathcal{L}_k$.

Now we apply to the sequence $\mathcal{L}_1 \supset \mathcal{L}_2 \supset \dots$ the p -adic analog of the lemma on the imbedded closed intervals of real analysis. The analog of this lemma holds for the topological spaces of much more general type — see e.g. the theorem in [16, Ch. 3, section 34, I], from which the p -adic case could be easily deduced. Thus, we conclude that the intersection of this sequence is not empty. That is, there exists $\mathbf{c}'' \in \mathbb{Z}_p^{r+2}$ which satisfies the assumptions of lemma 5.2. Yet then $d = 1$. $\square$

From here we deduce the following

5.3 Theorem. *Let $f \in \mathbb{Q}_p[x]$ be an integer-valued compatible and ergodic polynomial with $\deg f > 1$, and let there exists $r \in \mathbb{N}$ such that for each $k \in \mathbb{N}$ the*

linear complexity over the ring $\mathbb{Z}/p^k$ of the recurrence sequence $\{x_n\}$, defined by the recurrence relation $x_{n+1} \equiv f(x_n) \pmod{p^k}$, does not exceed r . In other words, let there exist $c^{(k)}, c_0^{(k)}, \dots, c_r^{(k)} \in \mathbb{Z}_p$ such that

$$c^{(k)} + \sum_{i=0}^r c_i^{(k)} x_{n+i} \equiv 0 \pmod{p^k} \quad (n = 0, 1, 2, \dots). \quad (\blacktriangleleft)$$

Then $\lim_{k \rightarrow \infty} c^{(k)} = \lim_{k \rightarrow \infty} c_1^{(k)} = \dots = \lim_{k \rightarrow \infty} c_r^{(k)} = 0$.

Proof. To start with, we note, that from the proofs of both lemma 5.2 and proposition 5.1 it follows that they remain true if we let k within their statements range over arbitrary infinite subset of $\mathbb{N}$.

Now for each $k \in \mathbb{N}$ choose (and fix) $c^{(k)}, c_0^{(k)}, c_1^{(k)}, \dots, c_r^{(k)} \in \mathbb{Z}_p^{(r+2)}$ satisfying $(\blacktriangleleft)$. Put $\mathbf{c}_k = (c^{(k)}, c_0^{(k)}, c_1^{(k)}, \dots, c_r^{(k)}) \in \mathbb{Z}_p^{(r+2)}$. In view of 5.1 then $\|\mathbf{c}_k\|_p < 1$ for all $k \in \mathbb{N}$. Denote $\mathcal{N} = \{k \in \mathbb{N} : \|\mathbf{c}_k\|_p > p^{-k}\}$. In other words, $k \notin \mathcal{N}$ iff $(\blacktriangleleft)$ is equivalent to a congruence $0 \equiv 0 \pmod{p^k}$.

It is obvious that if $\mathcal{N}$ is finite, then the conclusion of the theorem is true. Let $\mathcal{N}$ be infinite.

For $k \in \mathcal{N}$ put $\hat{\mathbf{c}}_k = \|\mathbf{c}_k\|_p \mathbf{c}_k$ and denote $\hat{\mathcal{N}}$ a set of all $m \in \mathbb{N}$ such that $p^k \|\mathbf{c}_k\|_p = p^m$ for a suitable $k \in \mathcal{N}$. In other words, we replace each $(\blacktriangleleft)$ with the equivalent system of congruences

$$\hat{c}^{(k)} + \sum_{i=0}^r \hat{c}_i^{(k)} x_{n+i} \equiv 0 \pmod{p^m} \quad (n = 0, 1, 2, \dots),$$

where $(\hat{c}^{(k)}, \hat{c}_0^{(k)}, \hat{c}_1^{(k)}, \dots, \hat{c}_r^{(k)}) = \hat{\mathbf{c}}_k$, $p^m = p^k \|\mathbf{c}_k\|_p$.

If the set $\hat{\mathcal{N}}$ is finite, the conclusion of the theorem is obviously true. If $\hat{\mathcal{N}}$ is infinite, then, since $\|\hat{\mathbf{c}}_k\|_p = 1$, in view of 5.1 and the note at the beginning of the proof we conclude that $\deg f = 1$. A contradiction. $\square$

In the statement of the theorem 5.3 we mention a notion of linear complexity of a sequence over a ring. This is commonly used (especially in cryptography) characteristic of a quality of a sequence. Lemma 5.2 in these terms asserts that the sequence $\{x_i = f(x_{i-1}) : i \in \mathbb{N}\}$ has infinite linear complexity over $\mathbb{Z}_p$, providing $f \in \mathbb{Q}_p[x]$ is integer-valued compatible ergodic polynomial of degree $d > 1$. This assertion could be slightly strengthened.

5.4 Corollary. *If $f \in \mathbb{Q}_p[x]$ is an integer-valued compatible ergodic polynomial of degree $d > 1$, then a recurrence sequence $\{x_n\}$, which satisfy recurrence relation $x_{n+1} = f(x_n)$, has infinite linear complexity over $\mathbb{Q}_p$.*

Proof. If for suitable $c, c_0, \dots, c_r \in \mathbb{Q}_p$, which are not 0 simultaneously, the equality $c + \sum_{j=0}^r c_j x_{n+j} = 0$ holds for all $n = 0, 1, 2, \dots$, then the equality $hc + \sum_{j=0}^r hc_j x_{n+j} = 0$ with $h = 1$, if $c, c_0, \dots, c_r \in \mathbb{Z}_p$, and $h = \|(c, c_0, \dots, c_r)\|_p$ otherwise, holds either. In view of compatibility of f the conclusion now follows from 5.2. $\square$

Note. The assumption $f \in \mathbb{Q}_p[x]$ within statements of 5.1–5.4 can not be omitted. For instance, let $p = 2$ and let

$$f(x) = 1 + x + 4(-1)^{1+x} = 1 + x + \sum_{j=0}^{\infty} (-1)^j 2^{j+2} \binom{x}{j}.$$

By the theorem 2.3, the integer-valued function f is compatible and ergodic. However, it is easy to see that the recurrence sequence $\{x_n \in \mathbb{Z}_2\}$ with recurrence relation $x_{n+1} = f(x_n)$ satisfy the relation $x_{n+2} = x_n + 2$, i.e., has linear complexity 2 over $\mathbb{Z}_2$.

We should notice that in this section we use the notion of linear complexity of a sequence over a ring in a somewhat broader sense than it is commonly used. More often the linear complexity of a sequence $\{x_n\}$ of elements of a commutative ring R is understood as the smallest $r > 0$ such that there exist $c_0, \dots, c_{r-1} \in R$ which satisfy simultaneously all equations $x_{n+r} = \sum_{j=0}^{r-1} c_j x_{n+j}$ for $n = 0, 1, 2, \dots$. We, in distinction from it, admit nonzero constant term, as well as relations where all coefficients are zero divisors (yet not all 0 simultaneously; in the assertion of 5.3 the latter, however, is not important). If R is a field, then both notions basically do not differ one from another: if a sequence satisfies a relation $c + \sum_{i=0}^r c_i x_{n+i} = 0$ with $c_r \neq 0$, then it satisfies the relation $x_{n+r+1} = c_r^{-1} c_0 x_n - \sum_{j=0}^{r-1} c_r^{-1} (c_j - c_{j+1}) x_{n+j+1}$. Our definition seems to us some more convenient for geometric interpretations, see above.

In other words, we have shown that, loosely speaking, nonlinear ergodic polynomial generators are absolutely nonlinear — the sequences they produce can not be implemented as linear recurrences over $\mathbb{Q}_p$. We do not discuss here what is the impact of these results on testing of the corresponding generators with the above mentioned statistical tests, leaving this issue to the forthcoming paper. We only note that they give some evidence that nonlinear congruential generators with integer-valued compatible ergodic polynomials over $\mathbb{Q}$ as state change functions in practice will pass the tests.

Properly restated analogs of 5.1–5.4 hold for composite $m = p_1^{k_1} \dots p_s^{k_s}$, which is a product of powers of distinct primes $p_1, \dots, p_s$, providing the transformation f preserves all congruences of the ring $\mathbb{Z}_{p_1} \times \dots \times \mathbb{Z}_{p_s}$. In connection with congruential generators modulo a composite m we also note that one can take f to be a function, defined on the set $\mathbb{N}_0$ of all nonnegative integer rationals, which takes values in $\mathbb{Z}$, preserves all congruences of the ring $\mathbb{Z}$ and which is ergodic as a function of integer p -adic variable for all $p \in \{p_1, \dots, p_s\}$. These functions may also be constructed with the use of the results of the paper.

For instance, such functions may be found in the class

$$\mathcal{B}_0 = \left\{ \sum_{i=0}^{\infty} a_i (x)_i : a_i \in \mathbb{Z}; i = 0, 1, 2, \dots \right\},$$

where, we recall, $(x)_i$ is i th descending factorial power of x : $(x)_0 = 1$, $(x)_i = x(x-1) \dots (x-i+1)$ for all $i = 1, 2, \dots$. It is obvious that $\mathcal{B}_0$ is a proper subclass of the class $\mathcal{B}$ (studied in section 4) for each prime p (the definition of $\mathcal{B}$, we recall, depends on p). Since $\mathcal{B}$ consists of functions, which preserve all congruences of the ring $\mathbb{Z}_p$, then each function g of $\mathcal{B}_0$ preserves all congruences of the ring $\mathbb{Z}$, that is, for each $a, b \in \mathbb{N}_0$ and each natural number $N > 1$ a congruence $a \equiv b \pmod{N}$ implies a congruence $g(a) \equiv g(b) \pmod{N}$. So as a state change function of a pseudorandom generator we can take, for instance,

$$f(x) = 1 + x + p_1^2 \dots p_s^2 g(x) \quad (g \in \mathcal{B}_0);$$

in view of 4.12 f is ergodic as a function of integer p_j -adic variable for all $j = 1, 2, \dots, s$. That is, f is transitive modulo p_j^k for all $k = 1, 2, \dots$ and for all $j =$

$1, 2, \dots, s$. Thus, f is transitive modulo each $p_1^{t_1} \cdots p_s^{t_s}$, for arbitrary $t_1, \dots, t_s \in \mathbb{N}$. In particular, f is transitive modulo m , and hence a pseudorandom number generator with state change function f and arbitrary initial state $x_0 \in \{0, 1, \dots, m-1\}$ produces a purely periodic sequence of period length m , and each number of $\{0, 1, \dots, m-1\}$ occurs at the period of this sequence exactly once.

Obviously, $\mathcal{B}_0$ contains all polynomials with rational integer coefficients, so if $g(x) \in \mathbb{Z}[x]$ is a polynomial of degree $d \geq 1$, then the performance of the corresponding pseudorandom generator is equivalent to d additions and d multiplications modulo m of integer rationals. Obviously, $\mathcal{B}_0$ consists not only of polynomials over $\mathbb{Z}$. It is not clear, however, whether it contains other ‘natural’ functions which admit relatively simple program implementation.

Moreover, if m is arbitrary, it is not clear enough, which functions should be considered as ‘natural’, and which should not. If by ‘natural’ functions one understands compositions of arithmetical operations (addition, subtraction, multiplication, division, raising to a positive integer power, exponentiation) then the functions of this kind could be constructed, for instance, with the use of 2.3, 2.4, and 4.9 combined with 4.11 and 4.12. So, theorems 2.3–2.4 imply that a polynomial $f(x) \in \mathbb{Q}[x]$ of a form

$$f(x) = 1 + x + \sum_{i=0}^d c_i p_1^{\lfloor \log_{p_1}(i+1) \rfloor + 1} \cdots p_s^{\lfloor \log_{p_s}(i+1) \rfloor + 1} \binom{x}{i},$$

for arbitrary $c_0, c_1, c_2 \dots \in \mathbb{Z}$ is transitive modulo arbitrary natural number $M > 1$, which is a product of powers of $\{p_1, \dots, p_s\}$; in particular $f(x)$ is transitive modulo m . Hence, the performance of the corresponding pseudorandom generator is equivalent to d multiplications, d additions, $d+1$ reductions some moduli and one division of integer rationals.

From the above formula it follows that, for instance, a polynomial $f(x) = 1 + x + \frac{5}{18}(x)_6$ is transitive modulo 10^k for all $k = 1, 2, \dots$. In a similar way, with the use of 2.5 and 4.11 (or 4.9 together with 4.11) one may construct generators which use exponentiations. For instance, a function $f(x) = 1 + x + 201^x$ (or, more generally, a function $f(x) = 1 + x + (1 + 200u(x))^{w(x)}$ with $u(x), v(x) \in \mathbb{Z}[x]$), as well as a function $f(x) = 1 + x + 201^{201^x}$ are transitive modulo 10^k for all $k = 1, 2, \dots$ (see 4.9 and 4.11); the same is true for the function $f(x) = 1 + x + 100 \cdot 11^x$ (see 2.5 and 4.11). Judging by the number of publications on inversive generators, taking a multiplicative inverse (or, generally, raising to negative powers) modulo m also should be considered as ‘natural’ operations. Generators of this kind also could be constructed with the use of results of the paper: for instance, taking $w(x) = -1, v(x) = x$ in the just mentioned example, one obtains a function $f(x) = 1 + x + (1 + 200x)^{-1}$, which is transitive modulo 10^k for all $k = 1, 2, \dots$.

We note that during the past decade there were intensive studies of such pseudorandom generators, as power generator ($f(x) = x^r, r \in \mathbb{N}$), exponential generator ($f(x) = a^x$), twice exponential generator ($f(x) = a^{b^x}$) and inversive generator ($f(x) = a + bx^{-1}$ or $f(x) = (a + bx)^{-1}$). The examples of generators, which are mentioned above in the section, and which use compositions of arithmetical operations, including exponentiation and raising to negative power, as we see, are somewhat distinct from the ones usually studied (by summand $1 + x$, at the first turn). These distinctions practically do not worsen the performance of the corresponding programs. However, these distinctions do not allow us to apply to the generators considered in this paper the results on already studied generators. It

would be very useful to study the possibility of such transfer, since in this area there are a lot of important results belonging to different authors (unfortunately, we could not present even a short survey here because of huge number of these).

At the same time, all the generators, introduced in this paper, are modulo given m equivalent to generators with recurrence relation $x_{n+1} \equiv f_m(x_n) \pmod{m}$, where $f_m(x) \in \mathbb{Q}[x]$ (this is an immediate consequence of p -adic Weierstrass theorem, for the latter see e.g. [3, Ch. 10, Theorem 1]). Hence, all the results, obtained in literature for so-called polynomial congruential generators, could be immediately applied to generators, considered in this paper, at least, under extra restriction $f_m(x) \in \mathbb{Z}[x]$.

We should note also, that a number of generators, studied in literature, concern a specific case, when m is a product of two distinct large primes. The results of the current paper are of little interest for this particular case, since with the use of these results one can construct generators, which are either equivalent modulo a prime divisor p of m to linear congruential generator, or involve some given in advance transitive modulo p polynomial of degree > 1 . The latter must be constructed beforehand and then ‘adjusted’ to make it transitive modulo some p^s , with s satisfying assumptions of 3.14, 4.1 or 4.9. The methods of such ‘adjustment’ we hope to publish in one of forthcoming papers, and here we restrict ourselves with an example. For instance, using these techniques, and choosing a transitive modulo 5 polynomial $1+3x^3$, it is possible to construct a polynomial $1-127x-152x^3+152x^5$, which is transitive modulo each 10^k , with arbitrary $k = 1, 2, \dots$.

So in view of these considerations, methods of construction of pseudorandom generators, developed in the paper, could give the best effect if applied to the case when m is a product of relatively small primes raised to relatively large powers. Thus the case $m = 2^s$ is a natural focus point, being the easiest for program implementations, since the reduction of a positive integer rational modulo 2^s is merely a truncation of all its 2-base expansion senior bits, starting with the s th one (our numbering of digits starts with 0). It is this case, which leads to the most natural (judging by program implementation) operations other than the above mentioned arithmetical ones, namely, to bitwise logical operations like XOR, OR, AND and other bitwise operations with nonnegative rational integer operands, represented as 2-base expansions. And, luckily, there is a complete description of measure-preserving (or ergodic) functions in this case — see section 2 of the paper.

The obtained results make it possible to construct pseudorandom number generators, which satisfy some requirements to performance, statistics or cryptographic security. This theme will be thoroughly studied in forthcoming papers. Here we briefly note only that application of equiprobable functions, which are also studied in the paper, as output functions of congruential generators with ergodic state change functions, allows us, preserving uniformity of distribution, to eliminate one more well known disadvantage of congruential generators, the so-called ‘low bit effect’. The latter demonstrates each sequence $\{x_n\}$, satisfying recurrence relation $x_{n+1} \equiv f(x_n) \pmod{2^k}$ with compatible $f: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2$: a sequence, composed of j th digits of each x_n , has a period length at most 2^{j+1} only. Methods of remedy will be also studied in one of the future papers.

REFERENCES

1. L. Kuipers and H. Niederreiter, *Uniform distribution of sequences*, John Wiley & Sons, N.Y.–London–Sidney–Toronto, 1974.

2. D. E. Knuth, *The art of computer programming*, Vol. 2: Seminumerical Algorithms (3rd edition), Addison-Wesley Publ. Co, 1998.
3. K. Mahler, *p-adic numbers and their functions*, (2nd edition), Cambridge Univ. Press, 1981.
4. R. C. Alperin R. C., *p-adic binomial coefficients mod P* , Amer. Math. Month. **92** (1985), no. 8, 576–578.
5. R. R. Hall, *On pseudo-polynomials*, Arch. Math. **18** (1971), 71–77.
6. N. Koblitz, *p-adic numbers, p-adic analysis, and zeta-functions*, Springer-Verlag, New York, etc., 1977.
7. G. Marsaglia, *Random numbers fall mainly in the planes*, Proc. Nat. Ac. Sci. USA **61** (1968), 25–28.
8. H. Lausch and W. Nöbauer, *Algebra of polynomials*, North-Holl. Publ. Co, Amsterdam, 1973.
9. H. K. Kaiser and W. Nöbauer, *Permutation polynomials in several variables over residue class rings*, J. Austral. Math. Soc. **A43** (1987), 171–175.
10. I. A. Yurov, *On p-adic functions which preserve Haar measure*, Matematicheskie Zametki **63** (1998), no. 6, 935–950. (Russian)
11. V. S. Anashin, *Uniformly distributed sequences of p-adic integers*, Matematicheskie Zametki **55** (1994), no. 2, 3–46 (Russian); English transl. in Mathematical Notes **55** (1994), no. 2, 109–133.
12. V. S. Anashin, *Uniformly distributed sequences over p-adic integers*, Number theoretic and algebraic methods in computer science (A. J. van der Poorten, I. Shparlinsky and H. G. Zimmer, eds.), Proceedings of the Int'l Conference (Moscow, June–July, 1993), World Scientific, 1995, pp. 1–18.
13. R. Rivest, *Permutation polynomials modulo 2^w* , Finite fields and appl. **7** (2001), no. 2, 287–292.
14. V. S. Anashin, *Uniformly distributed sequences over p-adic integers ...*, Number theoretic and algebraic methods in computer science, (Conference abstracts. Moscow, 29 June–2 July, 1993), Int'l Centre for Sci. and Tech. Information, Moscow, 1993, pp. 6 – 8.
15. M. V. Larin, *Transitive polynomial transformations of residue class rings*, Discrete Mathematics / Diskretnaya Matematika (to appear). (Russian)
16. K. Kuratowsky, *Topology*, Vol. 1, Academic Press, N.Y.–London, 1966.
17. V. S. Anashin, *Uniformly distributed sequences in computer algebra, or how to construct program generators of random numbers*, J. Math. Sci. **89** (1998), no. 4, Plenum Publishing Corp., New York, 1355 – 1390.

FACULTY FOR THE INFORMATION SECURITY. RUSSIAN STATE UNIVERSITY FOR THE HUMANITIES. MIUSSKAYA SQUARE, 6, MOSCOW 125267, RUSSIA.

E-mail address: vladimir@anashin.msk.su, anashin@rsuh.ru